

# Woodland Caribou In Ontario

## Background To A Policy

CA20N  
NR  
- 1989  
W53



Ministry of  
Natural  
Resources

Vincent G. Kerrio  
Minister



Digitized by the Internet Archive  
in 2022 with funding from  
University of Toronto



CA201  
NR  
-1989  
65

# Woodland Caribou In Ontario

## Background To A Policy

W.R. Darby

H.R. Timmermann

J.B. Snider

K.F. Abraham

R.A. Stefanski

C.A. Johnson

January, 1989



Ministry of  
Natural  
Resources

Vincent G. Kerrio  
Minister

ISBN 0-7729-4449-0

© 1989, Queen's Printer for Ontario

Printed in Ontario, Canada

Single copies of this publication are for sale from the addresses noted below.

Current publications of the Ontario Ministry of Natural Resources, and price lists, are obtainable through the Ministry of Natural Resources Public Information Centre, Room 1640, Whitney Block, 99 Wellesley Street West, Toronto, Ontario M7A 1W3 (personal shopping and mail orders).

And:

Personal shopping: Ontario Government Bookstore, Main Floor, 880 Bay Street, Toronto.

Mail Orders: MGS Publications Services Section, 5th Floor, 880 Bay Street, Toronto, Ontario M7A 1N8. Telephone 965-6015. Toll free long distance 1-800 268-7540, in Area Code 807 dial 0-Zenith 67200.

Cheques or money orders should be made payable to the Treasurer of Ontario, and payment must accompany order.

# EXECUTIVE SUMMARY

Woodland caribou are a true symbol of Ontario's northern heritage and wilderness. They once inhabited all of Ontario south to Lake Nipissing, but since European settlement their distribution has steadily receded northward to its present southern limit of approximately 50 degrees latitude. The distribution is still shrinking. A few exceptions are remnant herds on the islands and northern shore of Lake Superior.

The Ontario public is generally unaware caribou exist in the province. The tourism and scientific value of caribou is largely unrecognized. Native hunting of caribou occurs under Treaty rights whereas non-Native hunting of caribou was prohibited in 1929.

Management of the species and its habitat in Ontario has been minimal. This document presents background information to a policy for woodland caribou that will strengthen awareness and management efforts. It summarizes the status, biology, issues and management of Ontario caribou.

No comprehensive program for caribou inventory exists in Ontario. Provincial population estimates have been made twice, once in 1965 and again in this document. The present population is estimated to be approximately 15,000 animals.

Caribou take longer to reach sexual maturity than other members of the deer family and do not have multiple births. This low reproductive potential, their diet and adaptations to survive winter, allow them to occupy the tundra and mature boreal forest niche. Conversely, moose and white-tailed deer occupy the young deciduous or mixed forest niche.

Predation and loss of mature coniferous forest are serious threats to Ontario caribou, especially where moose, deer and wolf numbers increase to relatively high levels in caribou range after logging. Forestry operations, high predator numbers and a parasite carried by white-tailed deer can impact negatively on caribou. Poaching and excessive Native hunting can also threaten the resource.

Current Ontario issues are: lack of awareness of the caribou resource; the shrinking distribution of caribou; mitigation of forestry effects on caribou; re-evaluation of the sport hunting closure; the economic value of caribou to Native people through subsistence use versus trophy sport hunting; predator management; and caribou relocations.

# TABLE OF CONTENTS

	PAGE		PAGE
EXECUTIVE SUMMARY .....	1	2.5 Harvest and Other Uses .....	19
TABLE OF CONTENTS .....	2	2.5.1 Harvest .....	19
LIST OF FIGURES .....	3	2.5.2 Viewing .....	20
LIST OF TABLES .....	3	2.5.3 Research .....	20
1.0 INTRODUCTION .....	4	3.0 MANAGEMENT HISTORY .....	21
2.0 THE RESOURCE AND ITS HABITAT .....	5	4.0 SPECIFIC ISSUES .....	21
2.1 Taxonomy and Description .....	5	4.1 Awareness of the Resource .....	21
2.2 Distribution and Numbers .....	5	4.2 Mitigation of Forestry Effects of Caribou ...	21
2.2.1 Former Distribution .....	5	4.3 Relocation of Caribou .....	21
2.2.2 Current Distribution .....	5	4.4 Predator Management .....	22
2.2.3 Population Estimates .....	5	4.5 Harvest .....	22
2.3 Biology .....	8	5.0 LITERATURE CITED .....	23
2.3.1 Reproduction .....	8	6.0 APPENDICES .....	27
2.3.2 Mortality .....	10	6.1 Appendix 1 - Potential Woodland Caribou	
2.3.3 Behaviour .....	10	Viewing Areas in Ontario ....	27
2.4 Habitat Use .....	15	6.2 Appendix 2 - Summary of Native People's	
2.4.1 Hudson Bay Lowlands .....	15	Harvest of Ontario Woodland	
2.4.2 Boreal Coniferous Forest .....	15	Caribou by Community	
2.4.3 Great Lakes - St. Lawrence Forest .....	17	1960 to 1984 .....	28
2.4.4 Forestry and Caribou .....	17	6.3 Appendix 3 - Bibliography of Ontario Wood-	
		land Caribou Reports and	
		Publications .....	29



# LIST OF FIGURES

	Page
1. Distribution of woodland caribou in Ontario in 1900, 1950 and 1985 (deVos and Peterson, 1951; Darby and Duquette, 1985; OMNR, unpublished data) . . . . .	6
2. Relationship of the 1985 southern limit of continuous caribou distribution in Ontario to an abrupt northward decline in observed moose density of adjacent wildlife management units (Darby and Duquette, 1985; OMNR, unpublished data 1975 to 1984) . . . . .	11
3. Relationship of the 1985 southern limit of continuous caribou distribution to estimated density of timber wolves across Ontario (Kolenosky, 1981; Darby and Duquette, 1985) . . . . .	12
4. Relationship of the 1985 southern limit of continuous caribou distribution in Ontario to average lynx catch per trapper during the 1977-78 season (Quinn, 1984; Darby and Duquette, 1985) . . . . .	13
5. Relationship of the 1985 southern limit of continuous caribou distribution to the forest regions of Ontario (Simkin, 1965a; Rowe, 1972; Darby and Duquette, 1985) . . . . .	16

# LIST OF TABLES

	Page
1. Recent woodland caribou population estimates for Ontario Ministry of Natural Resources districts within current caribou range. Confidence intervals( $P < 0.10$ ) are shown where reported . . . . .	7
2. Reproductive rates of female caribou by age from nine North American herds (Bergerud, 1980) . . . . .	8
3. Percentage calves found in late winter surveys of five North American caribou herds (Bergerud, 1974a) . . . . .	9
4. Estimates of population density and percentage calves for woodland caribou herds derived from winter aerial transect surveys, 1959 to 1985. Confidence intervals ( $P < 0.10$ ) are shown where reported . . . . .	9
5. Estimates of annual Native harvest of woodland caribou for Ontario Ministry of Natural Resources districts . . . . .	19

# 1.0 INTRODUCTION

This document provides background information for the development of a provincial policy for woodland caribou in Ontario. It is intended to provide the public and professionals with a summary of the status, biology, issues and management of Ontario caribou. It is based on results of a comprehensive literature review and analysis of caribou data.

The document is organized into four principle sections: a summary of current knowledge about caribou; a brief history of caribou management efforts in Ontario; a synopsis of major issues relating to Ontario caribou; and an appendix of published and unpublished information.

Ontario has been managing caribou in the absence of a provincial policy. This lack of direction has hampered an organized approach to management. Public awareness of the resource is poor at a time when caribou are increasingly threatened by a variety of factors. A stimulus is needed to increase public and government awareness of the value of caribou. It is the purpose of this document to support a provincial policy that will provide for sound management of the resource.



## 2.0 THE RESOURCE AND ITS HABITAT

### 2.1 Taxonomy and Description

The caribou is a medium-sized member of the deer family, Cervidae. It is a ruminant of the Order Artiodactyla; in other words, a mammal with a four compartment stomach and cloven hooves (Banfield, 1974). All races of caribou and reindeer belong to one species, (*Rangifer tarandus*) (Banfield, 1961). The caribou of Ontario are all considered woodland caribou (*R. t. caribou*), one of five sub-species present in North America.

Woodland caribou are larger, darker and less migratory than barren-ground caribou (*R. t. groenlandicus*) with which people are more familiar. Coloration varies slightly with age, sex and season, but is usually dark brown with a white neck and ventral mane, white groin and rump. The legs are brown with narrow white 'socks' above the hooves. The trunk and head are slightly elongated, and the nose is large with a square muzzle. Adult bulls weigh 180-270 kg (400-600 lbs.), while adult cows weigh 90-140 kg (200-300 lbs.) (Bergerud, 1978). Generally, both sexes grow antlers. Bulls grow large massive antlers with erect spreading main beams, prominent 'bez' tines that are forward hand-like branches of the main beams, and palmate 'brow' tines that point forward over the nose. Cows usually have smaller antlers that are variable in shape, but some cows never grow antlers. Bulls shed their antlers from November through March, while pregnant cows retain their antlers until May or June after calving.

Woodland caribou have specialized adaptations to withstand the rigors of cold, northern winters. Their winter pelage is very dense with long, brittle guard hairs and a close, crinkly underfur. The hair is not hollow as commonly thought, but filled with tiny air bubbles. This makes the coat highly insulative and very buoyant. The muzzle is well furred, and the hairless portion restricted to a small oval on the upper lip. The ears and tail are short and heavily furred. The unusually large feet have long dewclaws and crescent-shaped hooves which facilitate travel over snow-covered or boggy ground (Banfield, 1974). This gives caribou a very distinctive crescentic, four-toed track. The hooves are longer in winter than in summer, with fur over the pads, providing good traction, insulation and ability to dig through the snow (crater) for food.

### 2.2 Distribution and Numbers

#### 2.2.1 Former Distribution

Woodland caribou once inhabited all of Ontario south to Lake Nipissing (Clarke, 1938; Snyder, 1938; Snyder *et al.*, 1942; deVos and Peterson, 1951). Their distribution steadily receded northward after European settlement in the late 1800's (deVos and Peterson, 1951; Cringan, 1956). It is still receding (Fig. 1).

During the late 1800's and early 1900's, the distribution of moose (*Alces alces*) and white-tailed deer (*Odocoileus virginianus*) expanded northward (Snyder, 1938; Peterson, 1955; Cumming and Walden, 1970). It is likely the decline of caribou was due to a combination of factors; hunting, fire, land clearing, logging, increased predation due to increased densities of moose and deer, disease caused by brain worm (*Parelaphostrongylus tenuis*) and human disturbance (Klein, 1968; Anderson, 1971, 1972; Bergerud, 1974a; Geist, 1978; Bergerud *et al.*, 1984a).

#### 2.2.2 Current Distribution

The southern limit of continuous caribou distribution in Ontario is presently a line of approximately 50 degrees latitude. South of this line there now remains only a few small scattered populations southeast of Geraldton, and on the islands, and northern shore of Lake Superior (Fig. 1).

#### 2.2.3 Population Estimates

No comprehensive inventory program for woodland caribou encompasses all the caribou range of Ontario. However, a provincial population estimate can be constructed from recent systematic aerial transect surveys of large portions of the present range, plus various localized surveys. This method yields a current population estimate of approximately 15,000 caribou (Table 1).

Provincial population estimates have been constructed in the past. Simkin (1965a) estimated the provincial population to be 13,000 in 1965 based on systematic aerial surveys that sampled portions of northern Ontario from 1959 to 1964. Cringan (1956) estimated 7,200 caribou existed in Ontario based on a review of literature and small localized surveys. deVos and Peterson (1951) estimated there were between 1,300 and 3,000 caribou in Ontario, but admitted poor coverage of the important Patricia areas.




Figure 1: Distribution of woodland caribou in Ontario in 1900, 1950, and 1985 (deVos and Peterson, 1951; Darby and Duquette, 1985; OMNR, unpublished data).



Table 1: Recent woodland caribou population estimates for Ontario Ministry of Natural Resources' districts within current caribou range. Confidence intervals ( $P < 0.10$ ) are shown where reported.

District	Population	District Estimate	Year	Reference
Cochrane	WMU 26	373 $\pm$ 345	1983	Armstrong (1983)
Geraldton	WMU 1D			
	Kowkash	2709	1978	Hamilton (1978)
	Fleming L.			Jackson (1981)
Hearst	Nagagami L.	22 <sup>a</sup>	1983	OMNR (unpubl. data, Hearst)
	Ebbs-Orkney Tp.			
Ignace	Seseganaga L.	7	1987	J. Carson (pers. comm)
Kapuskasing	WMU 25			Lucking (1979)
	Oke-Ford Tp.	80 <sup>a</sup>	1984	OMNR (unpubl. data, Moosonee)
Kenora	Umfreville L.			
	Snowshoe L.	37 <sup>a</sup>	1983	B. Ranta (pers. comm.)
	Eagle L.			
	Werner L. Rd.			
Moosonee	District-winter	4528 $\pm$ 1075	1981-4	Thompson (1986)
	Ft. Severn -			OMNR (unpubl. data, Moosonee)
	Man. - summer	4800 <sup>b</sup>	1986	
Nipigon	Onaman L.			
	Ogoki Res.			
	Armstrong	278 <sup>a</sup>	1978	(OMNR unpubl. data, Nipigon)
	Oboshkegan			
	Wabakimi L.			
Red Lake	WPLUP Area	570	1978	OMNR (1982a)
Sioux				
Lookout	WPLUP Area	1750	1978	OMNR (1982a)
Terrace Bay	Caramat			
	Coldwell Penn			
	Flanders Tp.	476 <sup>a</sup>	1986	OMNR (unpubl. data, Terrace Bay)
	Hagarty Rd.			
	Pic Is.			
	Slate Is.			
Wawa	L. Superior			
	Michipicoten Is.	52 <sup>a</sup>	1986	OMNR (unpubl. data, Wawa)
	Pukaskwa N. Pk.			
	Montreal Is.			
PROVINCIAL TOTAL		15682 <sup>c</sup>	1978-86	

a: These estimates are based on empirical data.

b: The relationship between summer and winter aggregations west of Ft. Severn is unclear. Caribou seen on the tundra in summer likely winter in Manitoba, hence are included here.

c: The provincial total is not a statistically based estimate; caution is required when comparing to previous provincial estimates.



The survey techniques, areas sampled and assumptions of these estimates differ among studies, and from those used to generate Table 1. Thus a direct comparison of estimates to determine population trend is not valid.

However, the similarity between Simkin's (1965a) estimate and the 1987 estimate in Table 1 is intriguing. It is important to note that no statistical confidence limits for the provincial estimates can be generated from available data.

## 2.3 Biology

### 2.3.1 Reproduction

Caribou take longer to mature than other cervids and do not have multiple births. Generally, female caribou become sexually mature at 2 1/2 years of age. However, a few breed as yearlings and some do not breed until 3 1/2 years or older (Bergerud, 1978). The average pregnancy rate of mature female caribou 2 1/2 years or older for seven different North American herds is 84 % (Table 2).

In the Northwest Territories, Dauphine (1976) found that barren-ground caribou cows with poor fat reserves did not breed until 3 1/2 years of age or older. Age of first conception is related to condition in most cervid species (Verme, 1967, 1969; Clutton-Brock et al., 1982; Saether and Haagenrud, 1983).

Two estimates of pregnancy rate exist for Ontario caribou. Firstly, Simkin (1965a) reported a pregnancy rate of 75 % for cows 1 1/2 years and older in the Patricia

region of northern Ontario. He also found that calves comprised 16 % of the winter population. Assuming yearlings comprise 16 % of the adult population, pregnancy rate is therefore 89 % for cows 2 1/2 years and older. Secondly, Gray (1978) estimated pregnancy rate of cows 1 1/2 years and older in the West Patricia region to be 63.5 % between 1960 and 1969. If we again assume yearlings comprise 16 % of the adult population, then pregnancy rate would be 76 % for cows 2 1/2 years and older.

A factor that compensates for delayed maturity and single births in caribou is an adult sex ratio favouring females. The mean percent females of adult caribou in eight North American populations reported by Bergerud (1980) was 61 %. The preponderance of females means that calves at birth comprise 27-30 % of the population (Bergerud, 1980). The only information on the sex ratio of adult Ontario caribou comes from a small herd inhabiting Pukaskwa National Park that averaged 60 % females over an eight year period (Bergerud, 1985).

Calf recruitment rates commonly obtained during late winter aerial surveys are important indicators of population growth potential. The proportion of calves in five North American caribou herds surveyed in late winter ranged from 10 % to 16 % (Table 3), and averaged 12.4 % (Bergerud, 1974a). The proportion of calves in Ontario herds surveyed in winter ranged from 0.0 to 22.5 % (Table 4) and averaged 12.8 %.

Table 2: Reproductive rates of female caribou by age from nine North American herds (Bergerud, 1980).

Herd Name	% Pregnant				Total <sup>a</sup>	% Parous <sup>b</sup> 2+
	Calf	1.5	2.5	3+		
Kaminuriak (NWT)	0(48) <sup>c</sup>	2(57)	48(69)	90(280)	81	82(343)
Beverly (NWT)	—	33(3)	50(16)	78(69)	73	—
Nelchina (AK)	0(24)	13(31)	61(46)	89(335)	86	86(3526)
W. Arctic (AK)	—	—	—	78(130)	78	92(517)
George R. (PQ)	—	100(1)	100(6)	100(15)	100	89(1279)
Spatsizi (BC)	—	—	—	—	—	80(3530)
Level Mt. (BC)	—	—	—	—	—	85(105)
Interior (NFLD)	—	—	—	—	81(21)	85(6657)
Porcupine (AK)	—	—	—	—	86(123)	—
TOTAL	0(72)	8(92)	55(137)	85(829)	84 <sup>a</sup>	86

a: Excludes calves and yearlings

b: Observations made on calving grounds during immediate post-calving period. Some segregation of parous and non-parous cows may have occurred. The sample excludes calves from the previous year.

c: Sample size.

Table 3: Percentage calves found in late winter surveys of five North American caribou herds (Bergerud, 1974a).

Location & Years	Number of herds surveyed	Percentage calves in herd	
		Range	Mean
Ontario 1960-64	4	15-18	16
NW Territories 1955-61	6	7-25	15
George R., Labrador 1957-63	7	7-17	11
Kaminuriak, NWT- Man., 1967-69	3	9-11	10
Mealy Mt., Labrador 1957-63	7	3-15	10

Table 4: Estimates of population density and percentage calves for woodland caribou herds in Ontario derived from winter aerial transect surveys, 1959 to 1985. Confidence intervals ( $P < 0.10$ ) are shown where reported.

Year	Survey Location	Area Sampled (km <sup>2</sup> )	Caribou Density (#/km <sup>2</sup> )	% Calves	Reference
1984-85	Cochrane	29500	.014	—	Dawson & Payne (1985)
1981-84	Moosonee District	221164	.020 ± .005	—	Thompson (1986)
1975-85	Pukaskwa Nat'l Pk	—	—	18.9	Bergerud (1985)
1976-81	L. Nipigon	32000	.006	21.0	Cumming & Beange (1987)
1979	W. Patricia	223500	.014 ± .004	11.6	Hamilton (1979a)
1979	W. Patricia <sup>a</sup>	9445	.047	—	Hamilton (1979b)
1976	Irregular L.	4300	.011	4.3	English (1976)
1974	Irregular L.	4300	.008	0.0	English (1974)
1973	Irregular L.	4300	.006	4.0	Busch (1973)
1972	Big Trout L.	14117	.011	—	Buss & Barbowski (1974)
1971	Severn-Winisk	21760	.042	11.1	Buss (1971a)
1971	Irregular L.	4300	.008	11.8	Buss (1971b)
1968	Irregular L.	4300	.012	22.5	Hagan (1968)
1967	Severn-Winisk	21760	.022	18.8	Simkin (1967)
1967	Cliff L.	1600	.023	—	Hansson (1967)
1966	Big Trout L.	14117	.017	—	Simkin (1966)
1965	Irregular L.	4300	.020	—	Simkin (1965b)
1963	Irregular L.	4300	.006	—	Armstrong(1963)
1959-64	N. Ontario	293690	.021	16.7	Simkin (1965a)

a: High density wintering areas only.

To demonstrate gross productivity of Ontario caribou (gross population recruitment at calving) it is useful to refer to a hypothetical example. Let us assume:

- (1) a starting population of 100 caribou during early May prior to calving;
- (2) an adult sex ratio of 39 males:61 females (Bergerud, 1980);
- (3) a recruitment rate of 0.128 to one year of age (i.e., net productivity) based on an average 12.8% calves observed in late winter aerial surveys (Table 4); and
- (4) a fecundity rate for cows 1 1/2 years and older equal to 0.75 (Simkin, 1965a).

In early May, prior to calving, the population of 100 caribou is comprised of: 13 calves, 34 bulls ( $87.2 \times 0.39$ ) and 53 adult cows ( $87.2 \times 0.61$ ). During May, 40 new calves are born ( $53 \times 0.75$ ). The total number of caribou in the population is now 140. The gross productivity is therefore 28.6% [ $40/(40 + 100)$ ]. Given the net productivity of 12.8%, or 14.7 calves per 100 adults ( $12.8 \times 100/87.2$ ), mortality in the first year of life is 63.3% [ $(40-14.7)/40$ ].

It is apparent from this example that calf mortality probably plays a major role in the population dynamics of Ontario's caribou as it does in many caribou herds across North America (Bergerud, 1980). It is important that any management of caribou in Ontario assess the magnitude and causes of calf mortality and attempt to alleviate this restriction to population growth.

There are examples of rapid population growth in caribou herds where calf survival is good due to the absence of predators and hunting. In Ontario, caribou introduced to Michipicoten Island in 1982 increased from 8 to 16 in three years with the observed rate of increase  $r=0.33$  (Gord Eason, OMNR, Wawa, pers. comm.). In Newfoundland, Brunette Island caribou increased at  $r=0.35$  from 1962 to 1967 (Bergerud, 1980). In Alaska: Adak Island caribou increased at  $r=0.25$  from 1962 to 1968; St. George and St. Paul Island caribou increased at  $r=0.24$  from 1911 to 1921, and; Kenai Peninsula caribou increased at  $r=0.27$  from 1966 to 1972 (Bergerud, 1980).

### 2.3.2 Mortality

Predation is probably the most important mortality factor affecting Ontario's caribou. It has been suggested by Simkin (1965a) and argued by Bergerud (1974a), that the decline of caribou in Ontario in the early 1900's was a result of increased wolf predation. Moose expanded their range northward from 1880 to 1910 (Peterson, 1955). White-tailed deer did not invade northwestern Ontario until 1897 (Snyder, 1938). The increased ungulate density allowed higher wolf numbers, increasing predation on caribou, and resulting in a decline of caribou in the

southern portion of their range. If we superimpose the present southern limit of continuous caribou distribution in Ontario on a map with the northern limit of high moose density, there is little overlap (Fig. 2). Figure 3 illustrates wolf densities across Ontario (Kolenosky, 1981); a gradient of high to low wolf density is apparent from south to north. Other significant predators of caribou are lynx (*Lynx canadensis*) (Bergerud, 1971) (Fig. 4), black bear (*Ursus canadensis*) and humans (see Section 2.5.1). Train collisions with caribou, common near the southern limits of distribution, can have a significant effect on small populations [e.g., 7 killed in Talbot township of a population of about 35 (OMNR, Hearst District unpublished data); 5 killed of about 25 in the Kowkash herd (Jackson, 1981)]. Automobile collisions with caribou are rare.

### 2.3.3 Behaviour

'Caribou', for many people, evokes images of vast herds that migrate long distances. Such dramatic behaviour appears to be a response to living in open habitats, and is not characteristic of the forest-dwelling populations of woodland caribou. For example, barren-ground caribou of the Northwest Territories, and woodland caribou of northern Quebec and central Newfoundland, live in open habitats throughout much of the year. They migrate hundreds of kilometers in spring, in large aggregations, to common calving areas in open habitats where predator densities are low. In summer they occupy open habitats where forage is nutrient-rich during the short but intense growing season. They rut in open habitats along treeline in autumn, then migrate back to forested habitats to winter.

In mountainous regions of British Columbia and the Yukon, woodland caribou undergo seasonal altitudinal migrations (Oosenbrug and Theberge, 1980; Bergerud *et al.*, 1984b). Calving on alpine tundra is considered an anti-predator strategy similar to that of calving on northern tundra (Bergerud *et al.*, 1984b). In both cases, caribou form large post-calving aggregations in open habitats, then migrate, and spend winter in smaller groups in forested habitat.

Some woodland caribou in Ontario occupy coastal tundra habitat from late winter through summer and early fall. In early June, 1986, the authors carried out an aerial reconnaissance survey along the Hudson Bay coast. A dispersed herd of more than 2,500 caribou was discovered near the coast opposite the Pen Islands, approximately 30 km east of the Manitoba border. Most of these caribou were cows and calves. Small groups of bulls were found scattered for at least 80 km into Manitoba.






Figure 2: Relationship of the 1985 southern limit of continuous caribou distribution in Ontario to an abrupt northward decline in observed moose density of adjacent wildlife management units (Darby and Duquette, 1985; OMNR unpublished data 1975 to 1984).




Figure 3: Relationship of the 1985 southern limit of continuous caribou distribution to estimated density of timber wolves across Ontario (Kolenosky, 1981; Darby and Duquette, 1985).




Figure 4: Relationship of the 1985 southern limit of continuous caribou distribution in Ontario to average lynx catch per trapper during the 1977-78 season (Quinn, 1984; Darby and Duquette, 1985).



Forest-dwelling woodland caribou have evolved other strategies to avoid predators during calving. At winter's end, cows disperse to secretive sites where concealment is provided by the forest, topography and waterbodies (Bergerud and Page, 1987). Caribou aggregations in open habitat may be analogous to this strategy because concealment of the individual may be provided by the constant movement and mixing of the herd (Estes, 1974; Bergerud, 1974b; Shoesmith, 1978).

In forest habitat at Reed Lake, northern Manitoba, Shoesmith and Storey (1977) found that woodland caribou cows commonly calve on islands or near mainland shorelines. In southeastern Manitoba, Darby and Pruitt (1984) found some caribou cows calved on islands while others calved on mainland locations. In Quebec and Labrador, Brown *et al.* (1986) found that almost all forest-dwelling cows calved on islands or on string bogs, usually returning to the same locations to calve in successive years.

In Ontario, Simkin (1965a) verified calving on islands in the Irregular Lake area. Nearly 100 of more than 200 caribou near Lake Nipigon move over ice to islands of the lake before calving, and remain there until after freeze-up when they return to mainland wintering areas (Cumming and Beange, 1987). In Pukaskwa National Park on the northeastern shore of Lake Superior, caribou calve near the coastline or on offshore islands (Bergerud, 1985). Caribou on Pic Island, Lake Superior, live in rugged terrain where they can seek altitudinal security if threatened by wolves (Ferguson, 1982). On the Slate Islands of Lake Superior, where large mammalian predators are absent, caribou exist at a density of seven to 11 per km<sup>2</sup>, the highest density in North America (Bergerud, 1980).

It is evident from this summary that caribou cows seek islands, shorelines, bogs or rugged topography to calve as a strategy for predator avoidance and escape during spring and summer. This is apparently necessary to protect young calves from predation.

Average group size of forest-dwelling caribou populations varies throughout the year. In winter, average group size was 4.1 in Alberta (Fuller and Keith, 1981), 4.6 at Reed Lake, Manitoba (Shoesmith and Storey, 1977), 5.5 at Aikens Lake, Manitoba (Darby and Pruitt, 1984) and 10.6 at Lac Joseph, Quebec (Brown *et al.*, 1986). In Ontario, mean winter group size was 2.9 at Pukaskwa National Park (Bergerud, 1985), 8.6 near Lake Nipigon (Cumming and Beange, 1987) and 9.1 in the Moosonee OMNR district (Thompson, 1986). During aerial caribou surveys

across northern Ontario from 1959 to 1964, mean winter group size was 10.5 (Simkin, 1965a).

Forest-dwelling woodland caribou are essentially solitary during late spring and summer. They do not form large post-calving aggregations characteristic of barren-ground caribou. In the Birch Mountains of northeastern Alberta, average group size was 1.2 in July (Fuller and Keith, 1981). In Manitoba during summer it was 1.7 at Reed Lake (Shoesmith, 1978) and 1.2 at Aikens Lake (Darby and Pruitt, 1984). In Ontario, it was 1.5 at Pukaskwa National Park (Bergerud, 1985).

Simkin (1965a) collected information on monthly trends in mean group size of caribou by interviewing Native people in northern Ontario from 1959 to 1963. They reported that group size increased in September, remained relatively high through winter, declined in April or May prior to calving, and was just over 1 from June through August. This same trend has been observed in other forest-dwelling populations by Shoesmith and Storey (1977), Fuller and Keith (1981), Darby and Pruitt (1984) and Brown *et al.* (1986).

Range size occupied by forest-dwelling woodland caribou is also less than that occupied by open habitat populations. At Reed Lake, Manitoba, the individual range occupied by 11 radio-collared caribou averaged 253 km<sup>2</sup> in winter, 107 km<sup>2</sup> in spring and 13 km<sup>2</sup> in summer (Shoesmith and Storey, 1977). At Aikens Lake, Manitoba, the summer and autumn range for an adult cow was 28 km<sup>2</sup> and 56 km<sup>2</sup>, respectively (Darby and Pruitt, 1984). In the Birch Mountains of northeastern Alberta, summer ranges for bulls and cows averaged 77 km<sup>2</sup> and 63 km<sup>2</sup> respectively; autumn ranges for bulls averaged only 23 km<sup>2</sup>, and; winter ranges averaged 335 km<sup>2</sup> for bulls and 137 km<sup>2</sup> for cows (Fuller and Keith, 1981).

Seasonal range for entire populations has also been reported for some forest-dwelling situations. At Aikens Lake, Manitoba, a population of 30-40 caribou occupied 175 to 190 km<sup>2</sup> during summer and 95 to 140 km<sup>2</sup> during winter over a two year period (Darby and Pruitt, 1984). In a 32,000 km<sup>2</sup> study area at Lake Nipigon, Ontario, Cumming and Beange (1987) found that a population of more than 200 caribou wintered in scattered groups; wintering areas totalled 2,824 km<sup>2</sup> but averaged 390 km<sup>2</sup>.

## 2.4 Habitat Use

### 2.4.1 Hudson Bay Lowlands

This is an area of flat topography and poor drainage extending 100 to 350 km inland from the Hudson Bay and James Bay coasts (Fig.5; cf. Rowe, 1972). It is comprised of two sub-types. Immediately adjacent to Hudson Bay is the Coastal Tundra Belt (Fig. 5), a narrow strip of tundra which is primarily treeless with some patches of stunted white spruce (*Picea glauca*) and black spruce (*P. mariana*) (Simkin, 1965a; Ahti and Hepburn, 1967; Rowe, 1972). Further inland is the Sub-Arctic Lichen Belt (Fig.5) characterized by open bogs and patterned fens interspersed with open black spruce-lichen woodlands and tamarack (*Larix laricina*) (Simkin, 1965a; Rowe, 1972). On river-bank levees, conifer and hardwood forests occur that are similar in composition and growth to the Boreal Coniferous Forest further south (Rowe, 1972). South of James Bay, conifer-lichen woodlands are scattered, and black spruce and tamarack swamps are common (Ahti and Hepburn, 1967).

There is a paucity of information on caribou in the Hudson Bay Lowlands. Movements of caribou in this area appear to be variable and complex (Simkin, 1965a). Some caribou calve and summer in the Coastal Tundra Belt, and winter in the Sub-Arctic Lichen Belt (Simkin, 1965a; K. Abraham, OMNR, Moosonee, Pers. comm.). Other caribou occupy the Sub-Arctic Lichen Belt year round (Simkin, 1965a).

Caribou that summer on the coast exhibit migratory and grouping behaviour similar to barren-ground caribou and woodland caribou that migrate between open and forested habitats (Kelsall, 1968; Bergerud, 1972; Dauphine et. al, 1975; Oosenbrug and Theberge, 1980). In spring, groups of pregnant cows migrate to the coastal tundra to calve, and other age/sex groups follow later. Post-calving aggregations of up to 4800 caribou occur during late spring and early summer (K. Abraham, Pers. comm.). During summer, most feeding activity occurs on relatively dry sites such as old beach ridges and frost-heaved areas where terrestrial lichens and the leaves of willow (*Salix* sp.) and bog birch (*Betula glandulosa*) are eaten. Cotton grass (*Eriophrum* sp.), sedges (*Carex* sp.) and bog bean (*Menyanthes trifoliata*) are important foods in wet areas (Simkin, 1965a). Post-calving aggregations break up in late July, and some movement away from the coast may occur. Some caribou rut near the treeline in late September and October (J. Thompson, OMNR, Moosonee, Pers. comm.).

Migration inland occurs in late fall, but it is not known if special migration routes or wintering areas exist. No

caribou have been found wintering on the Coastal Tundra (Simkin, 1965a; Thompson, 1986).

Conversely, caribou that occupy the Sub-Arctic Lichen Belt year-round are essentially solitary in summer (Simkin, 1965a). This is typical of woodland caribou that occupy the Boreal Coniferous Forest (see below). Important summer foods of these caribou are willow, terrestrial lichens, cotton grass, horsetail (*Equisetum* sp.), bog birch, bog bean, and bog ericoids (Simkin, 1965a).

Both coastal and inland caribou winter in groups in the Sub-Arctic Lichen Belt. Brokx (1965) reported that most caribou winter in the central Lowlands on raised bogs where terrestrial lichens are abundant on ombrotrophic deposits of peat. Lichen woodlands, lichen heaths, palsa bogs and sinuously patterned peatlands are not selected. Terrestrial lichens, the staple winter food, are obtained by cratering through the snow, but bog ericoids, grasses, sedges, arboreal lichens and bog birch are also consumed (Simkin, 1965a).

### 2.4.2 Boreal Coniferous Forest

Extending south of the Hudson Bay Lowlands is a vast area of Boreal Coniferous Forest (Rowe, 1972) which spans the breadth of Ontario, south to the English River in the west, Lake Superior in the middle and Lake Abitibi in the east (Fig. 5). From west to east there is a gradual change from forests of black spruce, jack pine (*Pinus banksiana*) and poplar (*Populus* sp.) on broken relief of Precambrian granite ridges and intervening muskegs, to extensive forests of black spruce and muskeg on clay flats.

Woodland caribou inhabiting Boreal Coniferous Forest usually form groups of less than 50. Gregarious in autumn, winter and early spring, they are essentially solitary in late spring and summer (Simkin, 1965a; Shoesmith and Storey, 1977; Fuller and Keith, 1981; Darby and Pruitt, 1984; Edmonds and Bloomfield, 1984). In general, caribou use of mature and over-mature pine (*Pinus* sp.) and spruce forest is high, whereas use of deciduous forest is low; open bogs, lakes and islands are preferred for foraging, escape and calving (Simkin, 1965a; Euler *et al.*, 1976; Shoesmith and Storey, 1977; Fuller and Keith, 1981; Darby and Pruitt, 1984; Edmonds and Bloomfield, 1984). While caribou can use a variety of habitat types, they exhibit strong seasonal preferences governed by forage availability, predators and snow conditions.

In spring and summer, caribou feed on forbs, deciduous leaves, lichens, fungi, grasses and sedges (Simkin, 1965a; Bergerud, 1972). Where these foods are widely abundant, caribou use a greater diversity of habitats in spring and summer than in winter (Fuller and Keith, 1981; Darby and Pruitt, 1984). Spring dispersal of cows and calves,




Figure 5: Relationship of the 1985 southern limit of continuous caribou distribution to the forest regions of Ontario (Simkin, 1965a; Rowe, 1972; Darby and Duguet, 1985).



and their use of open bogs, lakes and islands, may be anti-predator strategies (Simkin, 1965a; Shoesmith and Storey, 1977; Fuller and Keith, 1981). Similarly, caribou may use shorelines, open bogs and exposed ridges for relief from insects (Shoesmith and Storey, 1977; Edmonds and Bloomfield, 1984).

In autumn and winter caribou feed on arboreal and terrestrial lichens, sedges and bog ericoids; woody browse is not a dietary staple (Simkin, 1965a; Bergerud, 1972; Darby and Pruitt, 1984; Edmonds and Bloomfield, 1984). Frozen lakes and creeks are used for travel, escape, resting and drinking slush water (Stardom, 1975; Fuller and Keith, 1981; Darby and Pruitt, 1984; Edmonds and Bloomfield, 1984).

Although it has been argued that lichens are not essential for caribou survival (Bergerud, 1972; Euler *et al.*, 1976), they are a valuable winter food. During winter, caribou metabolic rate and protein requirements are reduced while carbohydrate demands are relatively high. Lichens, although low in protein, are rich in carbohydrates (Russell and Martell, 1984). Nutrients missing in lichens are contained in evergreen shrubs and graminoids consumed (Klein, 1982).

In some areas of Boreal Coniferous Forest woodland caribou may use traditional migration routes to move between summer and winter range (Stardom, 1977; Edmonds and Bloomfield, 1984), but fidelity to such routes is not strict. Stardom (1977) reported the Sasaginigak herd in eastern Manitoba followed the same migration route in three of four years. In other areas caribou do not show cohesive and unidirectional movements and traditional routes do not appear to exist (Shoesmith and Storey, 1977; Fuller and Keith, 1981; Cumming and Beange, 1987). Individual or herd movements may be as great as 84 km (Edmonds and Bloomfield, 1984), but distances of 10-40 km are more common (Stardom, 1977; Fuller and Keith, 1981). Some caribou use the same range year round (Fuller and Keith, 1981; Darby and Pruitt, 1984; Edmonds and Bloomfield, 1984).

#### 2.4.3 Great Lakes - St. Lawrence Forest

South of the Boreal Coniferous Forest, the Great Lakes-St. Lawrence Forest (Rowe, 1972) spans the breadth of Ontario, divided in two by Lake Superior (Fig. 5). It is a mixed forest of greater botanical diversity than its northern counterpart. Characterized by eastern white pine (*Pinus strobus*), red pine (*Pinus resinosa*), eastern hemlock (*Tsuga canadensis*) and yellow birch (*Betula alleghaniensis*), it also contains boreal species and such dominants as: eastern white cedar (*Thuja occidentalis*), sugar maple (*Acer*

*saccharum*), red maple (*A. rubrum*), red oak (*Quercus rubra*), basswood (*Tilia americana*), and white elm (*Ulmus americana*).

The only populations of woodland caribou presently inhabiting Great Lakes - St. Lawrence Forest in Ontario are small introduced herds on Montreal and Michipicoten Islands in Lake Superior (Fig. 1).

#### 2.4.4 Forestry and Caribou

Woodland caribou in closed habitat occupy old growth, boreal coniferous forest. Caribou do not use woody browse as a staple winter food. In contrast, moose and white-tailed deer thrive in early successional forest; deciduous woody browse is the major portion of their diet in winter.

The importance of mature and over-mature coniferous forest to caribou conflicts with the forestry practice of clearcutting for paper production. Clearcut logging alters caribou habitat in more ways than simple reversion to an early successional stage unfavourable to slow-growing lichens. Stand conversion from coniferous to deciduous species may occur, especially where poplar (*Populus* sp.) are common. This is due to rapid vegetational reproduction of most deciduous trees and shrubs. Lichens, fungi and some ericoids are replaced by vasculars. Woody browse often proliferates, encouraging the growth of moose or deer populations, but not caribou. Increased moose and deer numbers cause increased wolf (*Canis lupus*) densities which result in increased predation rates on adjacent caribou (Bergerud and Page, 1987). Also, road access to caribou habitat is provided, facilitating native subsistence hunting, illegal hunting, and travel of wolves (Bergerud *et al.*, 1984a).

Reasons for the steady northward recession of caribou distribution in Ontario since European settlement are not clear, but it appears that a combination of factors has been involved. Logging, land clearing, fire and human disturbance may have forced caribou northward, or onto unsuitable range where their numbers declined (cf. Klein, 1968; Geist, 1978). Also, local populations may have been extirpated by hunting and increased predation and disease resulting from the growth of adjacent moose and deer populations (Anderson, 1971; Bergerud *et al.*, 1984a).

Since 1960 there have been several cases in northern Ontario where small groups of caribou (10 to 40) disappeared or were displaced when logging occurred in their range. Usually, documentation and data relating to these cases are poor or non-existent, for example at McKay Lake near Geraldton (Fig. 1). In three cases however, there are sufficient data to provide insight to the problem (Darby and Duquette, 1986).

*Case 1: Fleming Lake, Geraldton District* - For many years woodland caribou were known to winter near Fleming Lake north of Geraldton (Fig. 1). Twenty-one caribou were observed in December, 1981; other caribou were likely present but not observed (Mark Sobchuk, OMNR, Fort Frances, Pers. comm.). Forest composition of the winter range (170 km<sup>2</sup>) was 61% mature and over-mature conifer (80 yrs or older), 11% immature conifer, 9% deciduous forest, 7% mixed forest, 8% muskeg and open land, and 4% water.

From 1980 to 1983, 1130 ha of mature and over-mature conifer were cut in a southern 7% of the winter range: 640 ha were clearcut, and 500 ha were modified with rows of uncut blocks 5 ha in size, spaced 200 m apart. The modified clearcut was intended to protect sandy soils from erosion, improve regeneration success and secondarily mitigate the effects of cutting on caribou (M. Sobchuk, Pers. comm.).

Nine aerial transect surveys (1.3 hr. each) were flown to monitor caribou distribution after cutting; one per month during December, January and February of 1981-82, 1982-83 and 1983-84. Results showed that caribou did not occupy any of the cuts. Instead, they were found in adjacent mature and over-mature jack pine north of the cuts where terrestrial lichens were abundant. Some caribou tracks were observed along the northern boundary of the cuts, but none were observed in the disturbed area to 1986 (M. Sobchuk, Pers. comm.).

No data are available on caribou reproductive or mortality rates for this area. There is no evidence of human harvest of caribou in this winter range despite frequent patrols by Conservation Officers. Moose density (0.12/km<sup>2</sup>) did not increase after cutting and white-tailed deer are not present. Wolf density is low but unquantified. However, wolf predation of caribou is not likely to have increased after cutting, because moose density did not increase.

*Case 2: Armstrong, Nipigon District* - Since 1975 approximately 100 caribou have summered on islands in Lake Nipigon and most wintered on the mainland, some near Armstrong 10 to 50 km away from the islands (Fig. 1; Bergerud and Butler, 1975; Cumming and Beange, 1987; OMNR, unpublished data). The mainland winter range (180 km<sup>2</sup>) is comprised of 35% mature and over-mature conifer, 26% immature conifer (mainly jack pine 40 to 79 yrs old), 11% deciduous forest, 17% mixed forest, 7% muskeg and open land, and 4% water. Most caribou activity in winter occurs on jack pine-spruce-lichen uplands. The community of Armstrong (population 500), a road and the Armstrong airport are located on north-western and northern boundaries of the winter range.

From 1974 to 1985, several contiguous clearcuts totalling 1140 ha were made in mature and over-mature conifer in a western 6% of the winter range. Caribou have not used

the cutovers since they were cleared, but continue to use adjacent uncut winter range (Blake Beange, OMNR, Nipigon, Pers. comm.). This was observed: on aerial transect surveys in 1983 and 1985 to monitor caribou winter distribution; on stratified random aerial quadrant surveys for moose in January 1976, 1979 and 1984, and; during casual observations of caribou and caribou tracks near the Armstrong airport adjacent to uncut winter range, but not in cutovers.

No data are available on caribou reproduction or mortality rates for this area. There is no evidence of human harvest of caribou in this winter range despite frequent patrols by Conservation Officers. Moose density immediately west of the winter range was 0.16/km<sup>2</sup> in 1976 and 0.24/km<sup>2</sup> in 1979. To the southwest it was 0.36/km<sup>2</sup> in 1979. White-tailed deer are not present in the area. Wolves are present but no information is available on their density. Caribou numbers appear unaffected by the increased moose density west of the winter range, so increased wolf predation of caribou does not appear to be implicated.

*Case 3: Cliff Lake, Dryden District* - Brousseau (1978) reported caribou disappeared from the Cliff Lake area northwest of Dryden (Fig. 1) after logging occurred in their winter range. Caribou had been known to exist there for many years. Six aerial transect surveys in winter 1966-67 showed the number of caribou exceeded 36 (Hansson, 1967). Brousseau (1978) described how the distribution and number of caribou subsequently receded as cutting occurred from 1968 to 1978. Four aerial transect surveys in March, 1978, showed approximately 12 caribou remained, wintering on uncut rocky jack pine ridges. Annual pellet group surveys (Brousseau, 1978) showed caribou density declined from  $0.86 \pm 0.35/\text{km}^2$  ( $P < 0.05$ ,  $n=318$  plots) in 1972 to zero in 1978 ( $n=320$  plots). Cutting in the area has continued to date. No caribou have been seen in Dryden district since 1978 (W. May, OMNR, Dryden, Pers. comm.).

The winter range calculated from MNR sightings of caribou and caribou tracks, 1968 to 1978 (Hansson, 1967; Brousseau, 1978), was 270 km<sup>2</sup>. Before cutting, 39% was mature and over-mature conifer, 15% was immature conifer, 9% was deciduous forest, 11% was mixed forest, 9% was muskeg and open land, and 17% was water. By 1978, 15% was clearcuts, 60 to 1600 ha in size, scattered among lakes in central and eastern portions (33%) of the winter range. A road bisected the winter range, and large cuts and a road surrounded its eastern and northern margins. Cutting has continued in the area to date. Brousseau (1978) suggested the decline of possible emigration of caribou was due to logging. He speculated the reasons may have been direct, through habitat destruction and disturbance, or indirect through increased wolf predation and illegal hunting.

However, it appears other factors were involved. The density of white-tailed deer was high in the Cliff Lake area from the 1960's to approximately 1975 (W. May, Pers. comm.). No quantitative estimates of deer density are available for the Cliff Lake area, but deer density in a wintering area 50 km south was  $8.6 \pm 2.3$  per km<sup>2</sup> in 1964. Moose density in the Cliff Lake area ( $0.18 \pm 0.07$  per km<sup>2</sup>,  $P < 0.10$ ) did not appear to increase during the period of caribou decline. There is no information on wolf density, but wolf-killed deer were commonly seen on the lakes. A high density of wolves may have increased predation rates on caribou. The meningeal worm may have been a mortality factor (Anderson, 1971) because deer densities were high for at least 5 years before the caribou started to decline. No evidence of human harvest of caribou was observed during patrols by Conservation Officers from 1972 to 1977, (Carl Hansson, OMNR, Dryden, Pers. comm.), even though Brousseau (1978) mentioned the possibility.

## 2.5 Harvest and Other Uses

### 2.5.1 Harvest

Caribou hunting, other than by Native people, has been prohibited in Ontario since 1929. No estimate of illegal harvest is available. Assessment of Native harvest has

been historically through trapper's meetings (Simkin, 1965a), and more recently through interviews and questionnaires completed during land use planning initiatives (Gray, 1978; Thompson, 1986).

Major subsistence hunting occurs by coastal village Natives in the Hudson Bay Lowlands (400-500 caribou per year) and by interior village Natives in the northern portion of the Boreal Coniferous Forest (e.g., WPLUP area: maximum 150 caribou per year, Gray, 1978). Lower levels of Native harvest occur in other interior areas (Table 5).

In interior areas, Natives harvest an estimated two to seven percent of local caribou populations (Gray, 1978; OMNR, 1982a; OMNR, 1982b). Coastal Native communities may harvest as much as 12% of wintering caribou (Thompson, 1986). Further, due to the clumped distribution of caribou, particularly in winter, and because of Native hunting methods, local populations may be threatened by overharvesting (Simkin, 1965a; Thompson, 1986). Gray (1978) reported that 35% of caribou seen by Native hunters were killed.

Table 5: Estimates of annual Native harvest of caribou for Ontario Ministry of Natural Resources districts.

OMNR District	Herd or WMU	Annual Harvest	Year	Reference
Cochrane	WMU 26	10-15	1982	OMNR (1982b)
Geraldton	WMU 1C	90	1982	OMNR (1982c)
Hearst	Nagagami L.	0		S. St. Jules,
	WMU 18B	20-25 <sup>a</sup>	1983-84	OMNR, Pers. comm.
Ignace		0		
Kapuskasing	WMU 25	0		
Kenora	WMU 2 & 6	0		
Moosonee	WMU 1A, 1B, 1D, 25, 26	400-500	1974-77 1981-83	Thompson (1986)
Nipigon	WMU 18A	20	1983	OMNR (1982d)
Red Lake	WMU 2, 1C	15-20	1982	OMNR (1982a)
Sioux Lookout	WMU 1C, 16A	55-60	1982	OMNR (1982a)
Terrace Bay		0		
Wawa		0		
Ontario Total		610-730 <sup>b</sup>		

a: one complete herd killed

b: 4 to 5% of estimated Ontario population (Table 1).



### 2.5.2 Viewing

The mystique and wilderness quality of caribou hold high value for tourism. The factors creating good viewing opportunities are: natural settings, relatively high probability of encountering caribou or at least caribou sign, and a pleasant interpretive experience.

Several areas of the province have moderate or high potential for viewing caribou (Appendix 1). The island populations of Lake Superior, particularly Pic Island and the Slate Islands, represent excellent viewing opportunities near a heavily travelled tourist corridor. Woodland Caribou Provincial Park, Wabakimi Lake Provincial Park, and the Lake Nipigon Islands may have viewing potential at certain locations and times. Wilderness areas such as Polar Bear Provincial Park, the Pen Islands area of the Hudson Bay coast, and Kesagami Lake Provincial Park have more caribou than the southern sites; however, access is more difficult and the viewing opportunities could only be effectively provided through carefully controlled aerial tours.

The existence of woodland caribou in the province would surprise many Ontarians. The spectacular aggregations that occur seasonally along the Hudson Bay coast would capture the imagination of even seasoned travellers. There are significant opportunities for tourism promotion using caribou, and for viewing caribou, that are presently not being utilized.

### 2.5.3 Research

The historical decline of caribou in some parts of their range has been linked to factors including overhunting, increased natural predation, habitat destruction, human disturbance and disease (Anderson, 1972; Bergerud, 1974a; Geist, 1978). Bergerud (1985) studied dispersion and anti-predator strategies of caribou along the northern shore of Lake Superior. Dispersal and movements of caribou near Lake Nipigon were reported by Cumming and Beange (1987). A study of the population status and movements of the Pen Island herd of caribou along the Hudson Bay Coast was initiated by OMNR staff, Moosonee District in 1987.

Brox (1965), Simkin (1965a) and Ahti and Hepburn (1967) conducted studies of caribou habitat in northern Ontario. Darby and Duquette (1986) reviewed three examples of caribou-forestry interaction in Ontario and suggested several mitigative techniques to reduce logging impact. Darby and Pruitt (1984) studied habitat use of a population of 30-40 woodland caribou in a study area straddling the Manitoba-Ontario border. Additional habitat related studies were conducted in the Armstrong-Lake Nipigon

area by Bergerud and Butler (1975), Paulsen (1976a, 1976b), Antoniak (1979), Walroth (1980, 1981) and Cumming and Beange (1987).

Bergerud and his co-workers have been investigating the ecology of two island populations of caribou in Lake Superior; the Slate Islands (Bergerud and Butler, 1974; Butler and Bergerud, 1978; Dalton, 1985) and Pic Island (Ferguson, 1982). Lankester and his students at Lakehead University have been studying diseases and parasites of woodland caribou since the mid 1970's. Their work includes the description of nematodes (Lankester, 1976; Lankester *et al.*, 1976; Lankester and Northcott, 1979) and gastrointestinal helminthes of caribou (Lankester and Fruetel, 1986; Fruetel, 1987).

Bergerud (1974a, 1980) contended that caribou did not decline during the late 1800's and early 1900's due to a lack of lichens caused by habitat destruction and disturbance (Hypothesis I), but by increased mortality from hunting and predation (Hypothesis II). He proposed a test of these hypotheses: "Introduce caribou to an island lacking predators and lichens but having deciduous growth, sedges, and evergreen shrubs. Later, introduce wolves (before forage is overutilized)" (Bergerud, 1974a:769). He contended that Hypotheses I would be supported by a decrease in the caribou population prior to wolf introduction; Hypothesis II by a rise in population prior to wolf introduction, then a decrease as wolf predation increases (caribou forage must still be adequate to distinguish from Hypotheses I).

Bergerud's proposed test had a bearing on planning for reintroductions of caribou to areas where they could establish self-sustaining populations. Consequently, OMNR staff relocated two groups of eight Slate Island caribou to Michipicoten and Montreal Islands in 1982 and 1984 respectively, to perform Bergerud's test. Both islands were considered free of predators. By March 1987, Michipicoten caribou had increased to an estimated 20 to 25 animals; by fall 1987, Montreal Island caribou were estimated at 13 animals (G. Eason, OMNR, Wawa, Pers. comm.). A third relocation to Bowman Island, Lake Superior, where both wolves and deer exist, occurred in 1985 (Timmermann, 1985). These caribou were monitored with the use of radio-collars. Information derived from these studies will be valuable in formulating management strategies for rebuilding caribou populations elsewhere.

## 3.0 MANAGEMENT HISTORY

Early historical records indicate that white-tailed deer management was initiated in Ontario in 1821, but caribou received scant attention until after the turn of the 20th century. Concern for dwindling numbers of caribou in the 1920's resulted in hunting by non-Natives being prohibited in 1929. The season has remained closed since that time and management activity, while minimal, has been directed toward conservation and maintenance of the species. In the past three decades, management work has centered on population surveys, mortality assessment and research. Native harvest remains unregulated. Prohibition of non-Native hunting is enforced through normal enforcement patrols by field staff.

Five reviews of the status of Ontario caribou have been written (deVos and Peterson, 1951; Cringan, 1956; Simkin, 1965a; Bergerud, 1980; Darby and Duquette, 1986). There have been three large scale inventories over portions of caribou range in Ontario (Simkin, 1965a; Hamilton 1978, 1979a; Thompson, 1986), and numerous small scale inventories (see Section 2.2.3). Harvest assessments are summarized in Section 2.5.1 and research activities in Section 2.5.3.

Habitat management for caribou in Ontario has only been attempted since 1975. An attempt to modify logging practices to protect caribou habitat northwest of Dryden did not prevent disappearance of the herd (Brousseau, 1978). This and other attempts near Geraldton that basically applied moose habitat guidelines to caribou ended in failure. Attempts which restricted logging to peripheral portions of caribou range had more success (Darby and Duquette, 1986; cf. Section 2.4.4). Recently, OMNR draft habitat management guidelines for caribou have been developed (Chinook Consulting Co., 1985). Other recent efforts include relocating caribou to areas of former range (see Section 2.5.3), establishing Woodland Caribou Provincial Park (OMNR, 1986a) and proposing a public viewing site in Slate Islands Provincial Park (OMNR, 1986b).

## 4.0 SPECIFIC ISSUES

### 4.1 Awareness of the Resource

Ontario woodland caribou presently have low intrinsic value in the public mind. The public is generally unaware that caribou inhabit the province. This lack of awareness is an issue.

### 4.2 Mitigation of Forestry Effects on Caribou

At Fleming Lake and Armstrong (Fig. 1, see Sec. 2.4.4) it appears logging caused displacement of caribou from peripheral portions of their range, yet range abandonment did not result and caribou numbers did not decline. At Cliff Lake the disappearance of caribou probably resulted from a combination of factors: habitat destruction, predation, and possibly disease involving meningeal worm (Anderson, 1971).

Cutting of most mature conifer in an area may leave caribou no option but emigration. Small patch cutting, on the other hand, does not mitigate caribou displacement and may increase access for hunters and wolves, may increase moose and deer densities, and consequently increase wolf numbers. In Alberta, forest-dwelling (vs. mountain) caribou did not feed in clearcuts larger than 2 ha and were known to cross a larger cut only once in four years (Edmonds and Bloomfield, 1984).

While the above information does not clearly identify the reasons for caribou displacement or decline, it does indicate the important need to develop techniques to mitigate the effects of forestry on caribou (cf. Darby and Duquette, 1986). In view of the recent expansion of logging into previously remote areas of northern Ontario, the shrinking distribution of caribou is an issue.

### 4.3 Relocation of Caribou

Small remnant populations of woodland caribou still exist along the northern shore of Lake Superior and on its associated islands (Euler *et al.*, 1976; Ferguson, 1982; Bergerud, 1985). Eason (1982, 1984) stated that caribou living near the Lake Superior shoreline near Wawa were in danger of extirpation because of their small numbers. He and others believe a good case can be made for protecting and increasing caribou numbers as well as determining factors responsible for their decline. This can be achieved by supplementing small existing populations and establishing new ones in areas of former range, where they could be self-sustaining, through relocation of animals. Candidate stocking sites should be carefully researched to ensure they were formerly occupied by caribou, suitable habitat conditions for caribou currently exist, predation is minimal and deer numbers are low.

Caribou intended for relocation can be obtained from populations that periodically exceed their carrying capacity. Slate Islands Provincial Park offers such an opportunity. In recent years, the estimated caribou population on the Slates has fluctuated from  $\pm 600$  in 1983-84 to  $\pm 300$  in 1985-86, on a land area of 36 km<sup>2</sup>.

Capture and handling practices must conform to requirements issued under the Animals for Research Act 1972. Legislation, policies and procedures outlined in the Provincial Parks Act, North Central Regional Park Research Plan, and the Slate Island Provincial Park Master Plan must be recognized and followed. Relocation projects of this nature are regarded by the Ministry of Environment as research, as defined in Regulation 293 of the Environmental Assessment Act. Regulation 293, Section 12, exempts research undertakings from provisions of the Act (L. Stevens, OMOE, pers. comm.). The capture and handling of caribou may be an issue.

## 4.4 Predator Management

Predation may significantly depress caribou numbers in certain cases. This is an important management concern where Native harvest is high or where habitat disturbance has caused an increase in moose or deer numbers. In special circumstances where predators are contributing disproportionately to total mortality of caribou, selective efforts can be applied to reduce numbers of specific predators. Control of moose and deer numbers may also be necessary. Predator control is likely to be a controversial issue.

## 4.5 Harvest

Native people's harvest of caribou is unregulated. In most of the Ontario caribou range this harvest does not appear excessive, but in some areas it may exceed sustained yield (Thompson, 1986).

Non-native harvest of caribou is presently prohibited in Ontario, but many of the conditions that required season closure in 1929 no longer exist. Evidence that Ontario caribou populations are below carrying capacity of the range (Ahti and Hepburn, 1967), better inventories, modern management techniques such as selective harvest (e.g., trophy bulls only in specific wildlife management units) and the opportunity to increase recreational and economic benefits to northern Ontario society suggest the sport hunting closure be re-evaluated.

Thus, two provincial issues related to caribou harvest exist:

- (1) re-evaluation of the rationale for the season closure for sport hunting in 1929 in light of information and experience acquired since that time; and
- (2) the economic value of caribou to Native people through subsistence use versus trophy sport hunting opportunities marketed by Native people.



## 5.0 LITERATURE CITED

- AHTI, T. and R.L. HEPBURN. 1967. Preliminary studies on woodland caribou range, especially on lichen stands, in Ontario. Ont. Dept. of Lands and Forests Res. Rep. (Wildl.) No. 74. Toronto: 134 p.
- ANDERSON, R.C. 1971. Neurological disease in reindeer (*Rangifer tarandus tarandus*) introduced in Ontario. Can. J. Zool. 49(2):159-166.
- . 1972. The ecological relationships of meningeal worm and native cervids in North America. J. Wildl. Dis. 8:304-310.
- ANTONIAK, K. 1979. Forest wintering areas of woodland caribou in the Lake Nipigon area. Unpubl. rep., School of Forestry, Lakehead U., Thunder Bay. 15 p.
- ARMSTRONG. 1963. Caribou survey. Ont. Dept. of Lands & Forests unpubl. rep., Sioux Lookout.
- ARMSTRONG, E.R. 1983. The 1982-83 aerial survey of moose and woodland caribou in W.M.U. #26., Ont. Min. Nat. Res. unpubl. rep., Cochrane, 44 p.
- BANFIELD, A.W.F. 1961. A revision of the reindeer and caribou, genus *Rangifer*. Nat. Mus. Can. Bull. 177. 137 p.
- . 1974. The mammals of Canada. U. of Toronto Press, Toronto. 438 p.
- BERGERUD, A.T. 1971. The population dynamics of Newfoundland caribou. Wildl. Monographs No. 25. 55p.
- . 1972. Food habits of Newfoundland caribou. J. Wildl. Manage. 36(3):913-923.
- . 1974a. Decline of caribou in North America following settlement. J. Wildl. Manage. 38(4):757-770.
- . 1974b. The role of the environment in the aggregation, movement and disturbance behaviour of caribou. In: Geist, V. and Walther, F. (eds.). The behaviour of the ungulates and its relation to management. Vol. 2 IUCN New Ser. Publ. 24, Morges, Switzerland: 552-584.
- . 1978. Caribou. In: Schmidt, J.L. and Gilbert, D.L. (eds.). Big Game of North America, Ecology and Management. Stackpole Books, Harrisburg, Penn.: 83-101.
- . 1980. A review of the population dynamics of caribou and wild reindeer in North America. In: Reimers, E., Gaare, E., and Skjenneberg, S. (eds.). Proceedings Second Inter. Reindeer/Caribou Symp., Direktoratet for vilt og ferskvannsfisk, Trondheim: 556-581.
- . 1985. Antipredator strategies of caribou: dispersion along shorelines. Can. J. Zool. 63(6):1324-1329.
- . and BUTLER, H.E. 1974. Wildlife investigations on the Slate Islands. Ont. Min. Nat. Res. unpub. rep., Terrace Bay, 46 p.
- . and BUTLER, H.E. 1975. Some rambling thoughts on caribou distribution and abundance in the Armstrong - Nipigon region, 1975. Ont. Min. Nat. Res. unpubl. rep., Thunder Bay. 53 p.
- . H.E. BUTLER and D.R. MILLER. 1984b. Anti-predator tactics of calving caribou: dispersion in mountains. Can. J. Zool. 62:1566-1575.
- . R.D. JAKIMCHUK and D.R. CARRUTHERS. 1984a. The buffalo of the north: caribou (*Rangifer tarandus*) and human developments. Arctic 37(1):7-22.
- . and R.E. PAGE. 1987. Displacement and dispersion of parturient caribou at calving as antipredator tactics. Can. J. Zool. 65:1597-1606.
- BROKX, P.A.J. 1965. The Hudson Bay lowland as caribou habitat. Unpublished M.Sc. thesis, U. of Guelph, Guelph. 224 p.
- BROUSSEAU, C. 1978. Trends in the woodland caribou (*Rangifer tarandus*) population in the Cliff Lake area of the Dryden district 1972-1978. Ont. Min. Nat. Res. unpubl. rep., Dryden. 25 p.
- BROWN, W.K., J. HUOT, P. LAMOTHE, S. LUTTICH, M. PARE, G. ST. MARTIN and J.B. THEBERGE. 1986. The distribution and movement patterns of four woodland caribou herds in Quebec and Labrador. In: Gunn, A., Miller, F.L. and Skjenneberg, S. (eds.). Proc. Fourth Inter. Reindeer/Caribou Symp., Rangifer Spec. Issue No. 1:43-49.
- BUSCH, D. 1973. Irregular Lake study area aerial caribou survey - 1973. Ont. Min. Nat. Res. unpubl. rep., Sioux Lookout, 6 p.
- BUSS, M.E. 1971a. Aerial survey for woodland caribou in the Severn-Winisk rivers area 1971. Ont. Dept. Lands & Forests unpubl. rep., Sioux Lookout, 11 p.
- . 1971b. Irregular Lake study area aerial caribou survey 1971. Ont. Dept. Lands & Forests unpubl. rep., Sioux Lookout, 4 p.
- . and J. BARBOWSKI. 1974. Winter habitat utilization by woodland caribou in the Hudson Bay lowlands. Ont. Dept. of Lands & Forests unpubl. rep., Sioux Lookout, 12 p.
- BUTLER, H.E. and A.T. BERGERUD. 1978. The unusual story of the Slate Islands caribou. Nature Canada 1:37-40.
- CHINOOK CONSULTING LTD. 1985. Habitat management guidelines for woodland caribou. Unpubl. rep. prepared for Ont. Min. Nat. Res., Toronto. 24 p.

- CLARKE, C.H.D. 1938. A study of the mammal population of the vicinity of Pancake Bay, Algoma District, Ontario. Natl. Mus. Can., Bull. 88, p.141-152.
- CLUTTON-BROCK, T.H., F.E. GUINNESS and S.D. ALBON. 1982. Red deer: behaviour and ecology of two sexes. In: Schaller G.B. (ed.). Wildlife and Behaviour Ecology Series, Univ. of Chicago Press. 378 p.
- CRINIGAN, A.T. 1956. Some aspects of the biology of caribou and a study of the woodland caribou range of the Slate Islands, Lake Superior, Ontario. Unpubl. M.A. thesis, U. of Toronto. 300 p.
- CUMMING, H.G. and D.B. BEANGE. 1987. Dispersion and movements of woodland caribou near Lake Nipigon, Ontario. J. Wildl. Manage. 51(1):69-79.
- . and F.A. WALDEN. 1970. The white-tailed deer in Ontario. Ont. Min. of Nat. Res., Toronto. 24 p.
- DALTON, W.J. 1985. Background report for the Slate Island Provincial Park management plan. Ont. Min. Nat. Res. unpubl. rep., Terrace Bay.
- DARBY, W.R. and L.S. DUQUETTE. 1986. Woodland caribou and forestry in Northern Ontario, Canada. In: Gunn, A., Miller, F.L. and Skjenneberg, S. (eds.). Proc. Fourth Inter. Reindeer/Caribou Symp., Rangifer Spec. Issue No. 1:87-93.
- . and W.O. PRUITT, JR. 1984. Habitat use, movements and grouping behaviour of woodland caribou (*Rangifer tarandus caribou*) in southeastern Manitoba. Can. Field-Nat. 98(2):184-190.
- DAUPHINE, T.G. JR. 1976. Biology of the Kaminuriak population of barren-ground caribou. Part 4., Can. Wildl. Serv. Rep. Ser. No. 38. 71 p.
- . F.W. ANDERKA, C.A. DROLET and D.T. McILVEEN. 1975. Distribution and movements of marked caribou in Ungava, June 1973 to 1974. Can. Wildl. Serv. Prog. Note No. 46. 19 p.
- DAWSON, F.N. and D.A. PAYNE. 1985. The 1984-85 aerial survey of moose and woodland caribou in W.M.U. 26. Ont. Min. Nat. Res. unpubl. rep., Cochrane. 17 p.
- deVOS, A. and R.L. PETERSON. 1951. A review of the status of woodland caribou (*Rangifer caribou*) in Ontario. J. Mamm. 32:329-337.
- EASON, G. 1982. Caribou management proposal, Wawa district. Ont. Min. Nat. Res. unpubl. rep., Wawa. 11 p.
- . 1984. Caribou transfer proposal, Montreal Island. Ont. Min. Nat. Res. unpubl. rep., Wawa. 18 p.
- EDMONDS, E.J. and M. BLOOMFIELD. 1984. A study of woodland caribou (*Rangifer tarandus caribou*) in west central Alberta 1979-1983. Alta. Energy and Nat. Res., Fish and Wildl. Div. unpubl. rep., Edmonton, Alta. 203 p.
- ENGLISH, G. 1974. Aerial caribou survey - 1974 Irregular Lake study area. Ont. Min. Nat. Res. unpubl. rep., Red Lake. 3 p.
- . 1976. Aerial caribou survey - 1976 Irregular Lake study area. Ont. Min. Nat. Res. unpubl. rep., Red Lake. 2 p.
- ERIKSSON, O. 1975. Silvicultural practices and reindeer grazing in northern Sweden. In: Luick, J.R., Lent, P.C., Klein, D.R. and White, R.G. (eds.). Proc. First Inter. Reindeer/Caribou Symp., U. of Alaska Biol. Papers Spec. Rep. No. 1:108-121.
- ESTES, R.D. 1974. Social organization of the African bovidae. In: Geist, V. and Walther, F. (eds.). The behaviour of ungulates and its relation to management. IUCN New Ser. Publ. 24, Morges, Switzerland:166-205.
- EULER, D.L., B. SNIDER and H.R. TIMMERMAN. 1976. Woodland caribou and plant communities on the Slate Islands, Lake Superior. Can. Field-Nat. 90(1):17-21.
- FERGUSON, S.H. 1982. Why are caribou on Pic Island? Unpubl. M.Sc. thesis, U. of Victoria. 170 p.
- FRUETEL, M.J. 1987. The biology of the gastrointestinal helminthes of woodland and barren-ground caribou (*Rangifer tarandus*). M.Sc. thesis, Lakehead University, Thunder Bay, Ontario. 114 p.
- FULLER, T.K. and L.B. KEITH. 1981. Woodland caribou population dynamics in northeastern Alberta. J. Wildl. Manage. 45(1):197-211.
- GEIST, V. 1978. Behaviour. In: Schmidt, J.L. and Gilbert, D.L. (eds.). Big Game of North America, Ecology and Management. Stackpole Books, Harrisburg, Penn. 283-296.
- GRAY, P.A. 1978. Native peoples' harvest of woodland caribou (*Rangifer tarandus caribou*) in the West Patricia planning area, 1960 to 1978. Ont. Min. Nat. Res. unpubl. rep., Toronto. 39 p.
- HAGAN, D. 1968. Aerial caribou survey - Irregular Lake study area, 1968. Ont. Dept. of Lands & Forests unpubl. rep., Cochenour. 6 p.

- HAMILTON, G.D. 1978. Aerial census of woodland caribou in the West Patricia planning area. West Patricia Land Use Plan Wildl. Tech. Rep. No.6, Ont. Min. Nat. Res. Cochenour. 14 p.
- . 1979a. Reassessment of some woodland caribou populations in the west Patricia planning area. West Patricia Land Use Plan Wildl. Tech. Rep. No.15, Ont. Min. Nat. Res., Cochenour. 9 p.
- . 1979b. A study of some woodland caribou wintering areas within the West Patricia planning area. West Patricia Land Use Plan Wildl. Tech. Rep. No.16, Ont. Min. Nat. Res., Toronto. 21 p.
- HANSSON, C. 1967. The present status of woodland caribou in the Kenora district, 1966-67. Ont. Dept. of Lands and Forests unpubl. rep., Kenora. 6 p.
- JACKSON, G. 1981. Woodland caribou: report prepared for Geraldton district land use plan. Ont. Min. Nat. Res. unpubl. rep., Geraldton. 13 p.
- KELSALL, J.P. 1968. The migratory barren-ground caribou of Canada. Can. Wildl. Serv. Monogr. No. 3. 340 p.
- KLEIN, D.R. 1968. The introduction, increase, and crash of reindeer on St. Mathew Island. J. Wildl. Manage. 32(2): 350-367.
- . 1982. Fire, lichens and caribou. J. Range. Manage. 35:390-395.
- KOLENOSKY, G.B. 1981. Status of wolves in Ontario. In: Carbyn, L.N. (ed.) Wolves in Canada and Alaska. Can. Wildl. Serv. Rep. No. 45.
- LANKESTER, M.W. 1976. A protostrongylid nematode of woodland caribou and implications in moose management. In: Hancock, J.A. and Mercer, W.E. (eds.), Proc. 12th North American Moose Conference and Workshop: 173-190.
- . V.J. CRICHTON and H.R. TIMMERMAN. 1976. A protostrongyloid nematode (Strongylida: Protostrongylidae) in woodland caribou (*Rangifer tarandus caribou*). Can. J. Zool. 54:680-684.
- . and M. FRUETEL. 1986. Pathogenic helminthes of woodland caribou. Paper presented at the Ontario Renewable Resources Research Seminar, January 30-31, 1986, Toronto. 16 p.
- . and NORTHCOTT. 1979. (*Elaphostrongylus cervi*) Cameron 1931 (Nematoda: Metastrongyloidea) in caribou (*Rangifer tarandus caribou*) of Newfoundland. Can. J. Zool. 57(7): 1384-1392.
- LUCKING, E. 1979. Woodland caribou: a compendium of sightings in Kapuskasing district. Ont. Min. Nat. Res. unpubl. rep., Kapuskasing. 5 p.
- OMNR. 1982a. West Patricia land use plan: proposed policy and optional plans. Ont. Min. Nat. Res., Toronto. 368 p.
- . 1982b. Cochrane district land use guidelines. Ont. Min. Nat. Res., Cochrane. 84 p.
- . 1982c. Geraldton district land use guidelines. Ont. Min. Nat. Res., Cochrane. 84 p.
- . 1982d. Nipigon district land use guidelines. Ont. Min. Nat. Res., Nipigon.
- . 1986a. Slate Islands Provincial Park Management Plan; preliminary draft. Ont. Min. Nat. Res. unpubl. rep., Terrace Bay. 27 p.
- . 1986b. Woodland caribou provincial park background information. Ont. Min. Nat. Res., Red Lake. 82 p.
- OLOSENBURG, S.M. and J.B. THEBERGE. 1980. Altitudinal movements and summer habitat preferences of woodland caribou in the Kluane ranges, Yukon Territory. Arctic 3(1):59-72.
- PAULSEN, H.E. 1976a. 1975-1976 caribou studies. In: Swanson, R.J. (ed.) Nipigon District Annual Fish and Wildlife Report 1975/76, Toronto:128-129.
- . 1976b. Winter range of the Armstrong caribou herd. In: Swanson, R.J. (ed.) Nipigon District Annual Fish and Wildlife Report 1975/76, Ont. Min. Nat. Res. unpubl. rep. Toronto:124-126.
- PETERSON, R.L. 1955. North American moose. U. of Toronto Press, Toronto. 280 p.
- QUINN, N. 1984. The lynx. Ont. Min. Nat. Res., Toronto. 24 p.
- ROSS, R.K. 1985. A survey of migratory birds harvested by Native people from the communities of Big Trout Lake, Kasaboneka Lake and Wapekaka Lake. Canadian Wildlife Service unpubl. rep., Ottawa. 17 p.
- ROWE, J.S. 1972. Forest Regions of Canada. Can. For. Serv. Publ. No. 1300. 172 p.
- RUSSELL, D.E. and A.M. MARTELL. 1984. Winter range ecology of caribou (*Rangifer tarandus*). In: Olson, R. et al. (eds.), Northern Ecology and Resource Management, U. of Alta. Press, Edmonton, Alta. 117-144.
- SAETHER, B.E. and H. HAAGENRUD. 1983 Life history of the moose: fecundity rates in relation to age and carcass weight. J. Mamm. 64(2):226-232.



- SHOESMITH, M.W. 1978. Social organization of wapiti and woodland caribou. Ph.D. dissertation, University of Manitoba, Winnipeg, Manitoba. 155 p.
- and D.R. STOREY. 1977. Movements and associated behaviour of woodland caribou in central Manitoba. Proc. Inter. Congr. Game Biol. 13:51-64.
- SIMKIN, D.W. 1965a. A preliminary report of the woodland caribou study in Ontario. Ont. Dept. of Lands and Forests Sect. Rep. (Wildl.) No. 59, Toronto, 76 p.
- 1965b. Woodland caribou in Ontario. Fish & Wildl. Rev.:2-8.
- 1966. Aerial survey for caribou in area northwest of Big Trout Lake, February 1966. Ont. Dept. Lands and Forests unpubl. res., Maple, 9 p.
- 1967. Aerial survey for caribou between Severn and Winisk rivers, March, 1967. Ont. Dept. of Lands and Forests unpubl. res. rep., Maple, 5 p.
- SNYDER, L.L. 1938. A faunal investigation of western Rainy River District Ontario. Trans. Roy. Can. Inst. Vol. 14, Part 1: 157-181. Contributions of the Royal Ontario Museum of Zoology, Toronto.
- E.B.S. LOGIER and T.B. KURATA. 1942. A faunal investigation of the Sault Ste. Marie region, Ontario. Contributions Royal Ont. Mus. Zool. No. 21.
- STARDOM, R.R.P. 1975. Woodland caribou and snow conditions in southeast Manitoba. In: Luick, J.R., Lent, P.C., Klein, D.R. and White, R.G. (eds.). Proceedings of the First International Reindeer/Caribou Symposium, University of Alaska Biological Papers Special Report Number 1. 436-461.
- 1977. A study of the winter ecology of woodland caribou, (*Rangifer tarandus caribou*) and comparison with some aspects of the winter ecology of moose, (*Alces alces andersoni*) and whitetail deer, (*Odocoileus virginianus dacotensis*) (Mammalia:Cervidae), in southeastern Manitoba. M.Sc. thesis, U. of Manitoba, Winnipeg, Man. 147 p.
- THOMPSON, J.E. 1986. Population and harvest surveys for the Ontario Hudson Bay Lowland caribou. Ont. Min. Nat. Res. unpubl. rep., Moosonee. 28 p.
- TIMMERMANN, H.R. 1985. Caribou relocation project, St. Ignace Island, Nipigon District, North Central Region. Ont. Min. Nat. Res., North Central Region unpubl. rep., Thunder Bay. 13 p.
- VERME, L.J. 1967. Influence of experimental diets on white-tailed deer reproduction. Michigan Conserv. Dept. Res. and Develop. Rep. No. 100. 15 p.
- 1969. Reproductive patterns of white-tailed deer related to nutritional plane. J. Wild. Manage. 33(4):881-887.
- WALROTH, R. 1980. Caribou field trip, Whitesand River area, September 25, 1979; winter observations of Armstrong caribou herd 1979-80; assessment of caribou lichen plots. In: Ont. Min. Nat. Res. Nipigon District Ann. Fish & Wildl. Rep. 1979/80, Nipigon. p. 169-175.
- 1981. Caribou pellet group and lichen *Cladonia* survey, Armstrong, May 1981. Ont. Min. Nat. Res. unpubl. rep., Nipigon. 16 p.

# 6.0 APPENDICIES

## 6.1 APPENDIX 1: Potential woodland caribou viewing areas in Ontario.

Location & OMNR District	Type Opportunity	Constraints on Viewing	Potential for Tourism
Kesagami L. Cochrane	Fly-in canoe or boat trip, Prov. P.,	Fly-in only, scattered caribou in summer	Low
Kattawagami-Detour L. Cochrane	Self-accessed by road - fall & spring migration	Road condition	Low
Michipicoten Is. Wawa	Charter boat, aircraft Prov. Park	Small herd, weather	Moderate
Montreal Is. Wawa	Charter boat telescope from Trans-Canada Hwy. Lookout Prov. Park	Small herd, weather	High
Pukaskwa Nat'l Pk., Wawa	Vehicle access Nat'l Park	Small herd	Moderate
Pic Islands Terrace Bay	Boat	Small herd, weather	High
Slate Islands Terrace Bay	Charter boat, aircraft, Prov. Park	Weather, access	High
Nipigon Islands Nipigon	Boat, self-access	Weather, scattered caribou, disturbance	Low
Wabakimi L. Nipigon	Winter, Prov. Park	Elusive herds, may require carefully regulated air tour	Low-Moderate
Woodland Caribou Park Red Lake	Prov. Park canoe, fly-in or air tour	Scattered caribou in summer, elusive herds in winter	Moderate-High
Polar Bear Prov. Park Moosonee	Fly-in, or air tour, Cree guide	Cost, few caribou carefully regulated air tour	Moderate
Pen Islands Moosonee	Wilderness tundra, fly-in Cree guide or air tour	Cost, summer only, carefully regulated, disturbance	High

## 6.2 APPENDIX 2: Summary of Native peoples' harvest of Ontario woodland caribou by community, 1960 to 1984.

	Community <sup>a, b</sup>												
	Coastal							Interior					
Year	1	2	3	4	5	6	7	8	9	10	11	12	Total
60-61	—	—	—	—	—	—	—	—	2	—	—	—	2
61-62	—	—	—	—	—	—	—	—	2	3	—	—	5
62-63	—	—	—	—	—	—	—	—	6	—	—	—	6
63-64	—			—	—	—	—	—	7	1	—	—	8
64-65	61	18		—	—	—	—	—	2	—	—	7	88
65-66	92	25		—	—	14	—	—	—	—	—	15	152
66-67	98	41		—	—	10	—	—	3	2	—	2	156
67-68	48	37		—	—	3	—	—	13	—	—	1	102
68-69	53	39		—	—	9	—	—	1	—	—	—	102
69-70	63	30		—	—	2	—	—	1	—	—	—	96
70-71	52	129		—	—	1	—	—	2	6	—	—	190
71-72	21	25		—	—	2	—	—	11	21	—	—	80
72-73	60	36		—	—	8	—	—	27	3	—	10	144
73-74	43	22		—	—	—	—	—	7	—	—	—	72
74-75	43	47	20	—	—	15	96	—	3	—	—	—	224
75-76	61	7	26	—	—	6	127	—	—	—	—	—	227
76-77	71	10	38	63	—	9	151	—	7	—	1	—	350
77-78	—	—	—	—	—	—	—	—	1	11	5	29	46
78-79	—	—	—	—	—	—	—	—	—	—	—	—	—
79-80	—	—	—	—	—	—	—	—	—	—	—	—	—
80-81	—	—	—	—	—	—	—	—	—	—	—	—	—
81-82	46	60	33	—	—	1	38	—	—	—	—	—	178
82-83	199	14	41	—	1	—	51	—	—	—	—	—	306
83-84	—	—	—	—	—	—	—	28	37	—	—	—	65

a: Communities are numbered as follows: 1 = Attawapiskat; 2 = Ft. Albany; 3 = Kasheshwan; 4 = Ft. Severn; 5 = Moose R.; 6 = Moosonee/Moose Factory; 7 = Winisk; 8 = Big Trout Lake; 9 = Kasabonika; 10 = Landsdowne House; 11 = Summer Beaver; 12 = Webequie.

b: References are: communities 1 to 7 - Thompson (1986); communities 8 & 9 - Ross (1985); communities 9 to 12 - Jackson (1981).



## 6.3 Appendix 3: Bibliography of Ontario Caribou Reports and Publications

- AHTI, T. 1959. Preliminary Survey of the Range of Woodland Caribou in Ontario. 10 p. (Presented at the Dorset Fur Advisory Comm. Meeting, Apr. 15-16, 1959). Library, Whitney Block, Toronto.
- AHTI, T. and R.L. HEPBURN. 1967. Preliminary Studies on Woodland Caribou Range, Especially on Lichen Stands in Ontario. 134 p. (Dept. of Lands and Forests, Research Report No. 74, March 1967). (2 copies) Library, Whitney Block, Toronto.
- ANDERSON, R.C. 1971. Neurological disease in reindeer (*Rangifer tarandus tarandus*) introduced in Ontario. *Can. J. Zool.* 49(2):159-166.
- ANDERSON, R.C. 1972. The ecological relationships of meningeal worm and native cervids in North America. (*In J. of Wildlife Diseases* 8: 304-310)
- ANONYMOUS. 1954-59. District Inventories of Moose, Caribou and Wolves. Library, Whitney Block, Toronto.
- ANONYMOUS. 1955. Trappers Meetings — Geraldton Forestry District. (*In Wildlife Management District Annual Reports 1954-55*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1956. Annual Report of the Patricia Central and Patricia West Trap-line Management Districts. 6 p. (*In Wildlife Management District Annual Reports 1955-56*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1957. Annual Report of the Patricia Central and Patricia West Trap-line Management Districts. 7 p. (*In Wildlife Management District Annual Reports 1956-57. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1957. Fish and Wildlife Report — Kapuskasing District. (*In Wildlife Management District Annual Reports 1956-57. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1958. Annual Report on moose, woodland caribou and wolves for the Geraldton District. Unpubl. Report, Geraldton District.
- ANONYMOUS. 1958. Annual Report of the Patricia Central, Patricia West and Sioux Lookout Trap-line Management Districts. 8 p. (*In Wildlife Management District Annual Reports 1957-58. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1958. Caribou — White River Management Report 1957-58. 1 p. (*In Wildlife Management District Annual Reports 1957-58. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1958. Kapuskasing Forestry District Report on the Status of Fur-bearing and Big Game Animals 1957-58. 4 p. (*In Wildlife Management District Annual Reports 1957-58. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1958. Report on the Status of Fur-bearing and Big Game Animals for the Year 1957-58; Cochrane District. 17 p. (*In Wildlife Management District Annual Reports 1957-58. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1959. Bird and Game Kills and Estimates — Moosonee Interim Reports. 5 p. (*In Wildlife Management District Annual Reports 1958-59. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1959. Caribou Inventory and Distribution Map — Cochrane Interim Reports. 2 p. (*In Wildlife Management District Annual Reports 1958-59. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1959. Kapuskasing Forestry District Report on the Status of Fur-bearing and Big Game Animals 1958-59. 5 p. (*In Wildlife Management District Annual Reports 1958-59. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1960. Big Game Kill — Sioux Lookout District Annual Fish and Wildlife Management Report 1959-60. 1 p. (*In Wildlife Management District Annual Reports 1959-60. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1960. Caribou Survey 1960 — Cochrane District Interim Reports. 7 p. (*In Wildlife Management District Annual Reports 1959-60. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1960. Cochrane District Caribou Inventory and Distribution Map. 2 p. (*In Wildlife Management District Annual Reports 1959-60. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1961. Caribou — Chapleau District. 1 p. (*In Wildlife Management District Annual Reports 1960-61. Part II*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1961. Game Kills and Estimates — Moosonee Interim Reports. 12 p. (*In Wildlife Management District Annual Reports 1960-61. Part I*). Library, Whitney Block, Toronto.

- ANONYMOUS. 1961. Kapuskasing Forestry District Report on the Status of Fur-bearing and Big Game Animals 1960-61. 5 p. (*In Wildlife Management District Annual Reports 1960-61. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1962. Game Kills and Estimates — Cochrane Interim Reports. 12 p. (*In Wildlife Management District Annual Reports 1961-62. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1964. Cochrane District Annual Report — Moosonee Area Game Kills. (*In Wildlife Management District Annual Reports 1963-64. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1965. Cochrane District Annual Report — Moosonee Area — Game Kills. (*In Wildlife Management District Annual Reports 1964-65. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1965. Moose, Caribou and Black Bear Killed by Means other than Legal. Library, Whitney Block, Toronto.
- ANONYMOUS. 1966. Chapleau Management Report 1965-66. 1 p. (*In Wildlife Management District Annual Reports 1965-66. Part II*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1966. Fish and Wildlife Management Report — Geraldton District. 1 p. (*In Wildlife Management District Annual Reports 1966-67. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1966. Geraldton District Annual Fish and Wildlife Management Report. (*In Wildlife Management District Annual Reports 1965-66. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1966. Kenora District Annual Fish and Wildlife Management Report 1965-66. (*In Wildlife Management District Annual Reports 1965-66. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1966. Moosonee Sub-Division Annual Fish and Wildlife Report and Interim Reports — Season 1965-66. (*In Wildlife Management District Annual Reports 1965-66. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1966. Sioux Lookout District Annual Fish and Wildlife Management Report 1965-66 — Wildlife Summary. (*In Wildlife Management District Annual Reports 1965-66. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1967. Chapleau Management Report 1966-67. 1 p. (*In Wildlife Management District Annual Reports 1966-67. Part II*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1967. Kenora District Annual Fish and Wildlife Management Report 1966-67. 1 p. (*In Wildlife Management District Annual Reports 1966-67. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1967. Sioux Lookout District Annual Fish and Wildlife Management Report 1966-67 — Big Game Kill. 2 p. (*In Wildlife Management District Annual Reports 1966-67. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1968. Kenora District Annual Fish and Wildlife Management Report 1967-68. 1 p. (*In Wildlife Management District Annual Reports 1967-68. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1968. Moosonee Sub-Division Annual Fish and Wildlife Report and Interim Reports — Season 1967-1968. (*In Wildlife Management District Annual Reports 1967-68. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1968. Sioux Lookout District Annual Fish and Wildlife Management Report 1967-68 — Wildlife Summary. (*In Wildlife Management District Annual Reports 1967-1968. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1968. White River District Fish and Wildlife Management Report 1967-68 — Big Game. (*In Wildlife Management District Annual Reports 1967-68. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1969. Kenora District Annual Fish and Wildlife Management Report 1968/69 — Big Game. 1 p. (*In Wildlife Management District Annual Reports 1968-69. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1969. Kesagami Lake Moose and Caribou Survey. 1 p. (*In Cochrane District Annual Fish and Wildlife Management Report 1968/69. p. 28*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1969. Moosonee Sub-Division Annual Fish and Wildlife Report and Interim Reports — Season 1968-1969. (*In Wildlife Management District Annual Reports 1968-69. Part III*). Library, Whitney Block, Toronto.

- ANONYMOUS. 1969. Sioux Lookout District Annual Fish and Wildlife Management Report 1968-69 — Big Game Kill. 2 p. (*In Wildlife Management District Annual Reports 1968-69. Part III*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1970. Fish and Wildlife Management Report 1969-70 — White River District. 39 p. (*In Wildlife Management District Annual Reports 1969-70. Part II*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1970. Kenora District Annual Fish and Wildlife Management Report 1969-70 — Big Game. 1 p. (*In Wildlife Management District Annual Reports 1969-70. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1970. Moosonee Sub-Division Annual Fish and Wildlife Report and Interim Reports — Season 1969-1970. (*In Wildlife Management District Annual Reports 1969-70. Part II*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1971. Moosonee Sub-Division Annual Fish and Wildlife Report and Interim Reports — Season 1970-1971. (*In Wildlife Management District Annual Reports 1970-71. Part II*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1971. Patricia East Spring Trapper's Meetings June 1970. 7 p. (*In Cochrane District Annual Fish and Wildlife Management Report 1970/71 and In Wildlife Management District Annual Reports 1969-70. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1971. Patricia East Spring Trapper's Meetings June 1971. 7 p. (*In Wildlife Management District Annual Reports 1970-71. Part I*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1972. Moosonee Sub-Division Annual Fish and Wildlife Report and Interim Reports — Season 1971/72. (*In Wildlife Management District Annual Reports 1971-72*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1976. Caribou Droppings. 1 p. (*In Nipigon District Annual Fish and Wildlife Report 1975/76. Ed. R.J. Swanson. p. 130*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1976. Winter Observations of Armstrong Caribou Herd. 1 p. (*In Nipigon District Annual Fish and Wildlife Rept. 1975/76. Ed. R.J. Swanson. p. 127*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1978. Caribou Plots — Supplement — Armstrong Airport. 1 p. (*In Nipigon District Annual Fish and Wildlife Report 1977-78. Ed. P. Burkhardt. p. 140*). Library, Whitney Block, Toronto.
- ANONYMOUS. 1978. Aerial Census Of Woodland Caribou In The West Patricia Planning Area, West Patricia Land Use Plan, Wildlife Technical Report No. 6, Kenora.
- ANONYMOUS. 1978. West Patricia Land Use Plan; Faunal Species List. 24 p. Library, Whitney Block, Toronto.
- ANONYMOUS. 1980. Woodland Caribou Collaring and Tagging, Onaman Lake. p. 165-68 (*In OMNR Nipigon District Annual Fish and Wildlife Report 1979*).
- ANONYMOUS. 1981. Timmins District Land Use Plan; Background Information. 153 p. (2 copies) Library, Whitney Block, Toronto.
- ANONYMOUS. 1981. West Patricia Land Use Plan; Background Information. 68 p. (3 copies) Library, Whitney Block, Toronto.
- ANONYMOUS. 1982. Caribou: Bibliography Selected From Wildlife Review 1935 — Mar. 1982. 97 p. Library, Whitney Block, Toronto.
- ANONYMOUS. 1982. Terrace Bay Land Use Plan; Background Information. 80 p. Library, Whitney Block, Toronto.
- ANONYMOUS. 1982. Wawa District Land Use Plan; Proposed Policy and Planning Options. Library, Whitney Block, Toronto.
- ANONYMOUS. 1982. Wawa District, Northeastern Region; District Land Use Planning Program; Background Information. 161 p. Library, Whitney Block, Toronto.
- ANONYMOUS. 1982. West Patricia Land Use Plan; Proposed Policy and Optional Plans. 368 p. (2 copies) Library, Whitney Block, Toronto.
- ANONYMOUS. 1983. Smoothrock Lake Caribou Tagging. p. 44-45 (P. Burkhardt (ed.) Nipigon District Annual Fish and Wildlife Report 1983).
- ANONYMOUS. 1983. Terrace Bay District Land Use Guidelines. 80 p. Library, Whitney Block, Toronto.
- ANONYMOUS. 1983. Thunder Bay District Land Use Guidelines. 108 p. Library, Whitney Block, Toronto.
- ANONYMOUS. 1983. Thunder Bay Land Use Plan; Background Information. 119 p. Library, Whitney Block, Toronto.



- ANONYMOUS. 1983. Timmins District Land Use Guidelines. 82 p. (2 copies) Library, Whitney Block, Toronto.
- ANONYMOUS. 1983. Wawa District; Land Use Guidelines. 80 p. Library, Whitney Block, Toronto.
- ANONYMOUS. Undated. Big Game Hunter Survey Report, 1968-69. Library, Whitney Block, Toronto.
- ANONYMOUS. Undated. Timmins District Land Use Plan; Proposed Policy and Planning Options. 97 p. (2 copies) Library, Whitney Block, Toronto.
- ANTONIAK, K. 1979. Forest Wintering Areas of Woodland Caribou in the Lake Nipigon Area. 15 p. (*In* Unpubl. Summer Progress Report). School of Forestry, Lakehead University, Thunder Bay, Ontario.
- ARMSTRONG, A.E. 1963. Caribou Survey. Ontario Department of Lands and Forests Unpubl. Report, Sioux Lookout District.
- ARMSTRONG, G.C. 1949. A General Outline Describing the Status of the Game and Predatory Animal in the Forest District of Port Arthur -1949. 11 p. Library, Whitney Block, Toronto.
- ARMSTRONG, E.R. 1980. The 1979-80 Aerial Inventory of Moose and Caribou in Wildlife Management Unit #26. 45 p. Library, Whitney Block, Toronto.
- ARMSTRONG, E.R. 1983. The 1982-83 Aerial Survey of Moose and Woodland caribou in Wildlife Management Unit #26. Ont. Min. Nat. Res. unpubl. rep., Cochrane District.
- BEANGE, D.B. 1982. Woodland Caribou Tagging Smoothrock Lake. p. 196-200. (*In* P. Burkhardt (ed.) OMNR Nipigon District Annual Fish and Wildlife Report 1983-83.
- BERGERUD, A.T. 1974. Decline of Caribou in North America following Settlement. (*In* J. Wildlife Management 38(4): 757-770).
- BERGERUD, A.T. 1978. Caribou. p. 83-101. (*In* J. Schmidt and Gilbert (eds) Big Game of North America). Stackpole Books.
- BERGERUD, A.T. 1979. Status of Rangifer in Canada. Woodland Caribou (*Rangifer tarandus caribou*). p. 748-753. (*In* E. Reimers, E. Gaare and S. Skjenneberg (eds.) Proc. 2nd Int. Reindeer/Caribou Symp. Roros, Norway 1979).
- BERGERUD, A.T. 1980. A Review of the Population Dynamics of Caribou and Wild Reindeer in North America. p. 556-581. (*In* E. Reimers, E. Gaare and S. Skjenneberg (eds.) Proc. 2nd Int. Reindeer/Caribou Symp. Roros, Norway 1979).
- BERGERUD, A.T. 1985. Antipredator Strategies of Caribou: Dispersion Along Shorelines. p. 1324-1329. (*In* Canadian J. Zool. 63).
- BERGERUD, A.T. and H.E. BUTLER. 1974. Wildlife Investigations on the Slate Islands. p. 46. (*In* Unpubl. Report OMNR Terrace Bay District.
- BERGERUD, A.T. and H.E. BUTLER. 1975. Some rambling thoughts on caribou distribution and abundance in the Armstrong-Nipigon region. 53 p. Unpubl. Report, Nipigon District.
- BERGERUD, A.T. and W.R. WYETT. 1976. Faunal Investigations on the Proposed Pukaskwa National Park. p. 63. (*In* Canadian National Parks, No. G-289).
- BERGERUD, A.T. and W.R. WYETT. 1976. Faunal Investigations in the proposed Pukaskwa National Park. p. 51. (*In* Unpubl. Report Dept. National Park Service, Pukaskwa National Park, Marathon, Ontario).
- BISHOP, C.A. 1974. The Northern Ojibwa and the Fur Trade: An Historical and ecological study. Holt, Rinehart and Winston of Canada, Ltd., Montreal.
- BROUSSEAU, C. 1979. Trends in the Woodland Caribou (*Rangifer tarandus*) Population in the Cliff Lake Area of the Dryden District. 25 p. Library, Whitney Block, Toronto.
- BUBENIK, A.B. 1975. Significance of Antlers in the Social Life of Barren Ground Caribou. 25 p. (*In* Proceedings of the First International Reindeer and Caribou Symposium, Aug. 9-11, 1972, University of Alaska, Fairbanks, Alaska). Library, Whitney Block, Toronto.
- BUBENIK, A.B. 1975. Taxonomic Value of Antlers in Genus *Rangifer*, H. Smith. 23 p. (*In* Proceedings of the First International Reindeer and Caribou Symposium, Aug. 9-11, 1972, University of Alaska, Fairbanks, Alaska). Library, Whitney Block, Toronto.
- BUSCH, D. 1973. Irregular Lake Study Area Aerial Caribou Survey 1973. Unpubl. Report, Sioux Lookout District. 6 p.
- BUSS, M.E. 1971. Aerial Survey for Woodland Caribou in the Severn-Winisk Rivers Area. Ontario Department of Lands and Forests, Unpubl. Report, Sioux Lookout District. 11p.
- BUSS, M.E. 1971. Irregular Lake Study Area Aerial Caribou Survey — 1971. Ontario Department of Lands and Forests, Sioux Lookout, 4 p.

- BUSS, M.E. 1972. Aerial Survey for Woodland Caribou Northwest of Big Trout Lake. Ontario Department of Lands and Forests, Unpubl. Report, Sioux Lookout District.
- BUSS, M.E. 1974. Caribou — Symbol of the Far North. 4 p. (Fish and Wildl. Review Vol 13(3): 3-6). (2 copies) Library, Whitney Block, Toronto.
- BUSS, M.E. and J.M. BARBOWSKI. 1974. Winter Habitat Utilization by Woodland Caribou in the Hudson Bay Lowlands. Ontario Department of Lands and Forests, Unpubl. Report, 12 p. Library, Whitney Block, Toronto.
- BUTLER, H.E. and A.T. BERGERUD. 1978. The Unusual Story of the Slate Islands Caribou. p. 37-40. (*In* Nature Canada January to March).
- BUTLIN, A. 1986. Protecting Lake Superior Caribou. 2 p. (*In* Aski, June 1986). Library, Whitney Block, Toronto.
- CHINOOK CONSULTING LTD. 1985. Habitat Management Guidelines for Woodland Caribou. Unpubl. Report for Ont. Min. Nat. Res., Toronto. 24 p.
- CLARKE, C.H.D. 1938. A study of the mammal population of the vicinity of Pancake Bay, Algoma District, Ontario. Natl. Mus. Can. Bull. 88, p. 141-152.
- CLARKE, C.H.D. 1947. Remarks on the Woodland Caribou, 1947. 2 p. (Ontario Department of Lands and Forests, 1947). Library, Whitney Block, Toronto.
- CLARKE, C.H.D. 1948. Caribou on the Islands of Lake Nipigon. (*In* Resources Management Report). Ontario Department of Lands and Forests, Toronto.
- CRAIG, M.H. 1961. Aerial Caribou Survey — Kapuskasing District. 6 p. (*In* Wildlife Management District Annual Reports 1960-61. Part I). Library, Whitney Block, Toronto.
- CRINGAN, A.T. 1948. Report upon Forest Conditions of Burnt Island in Connection with Woodland Caribou Range Requirements, 1948. 4 p. Library, Whitney Block, Toronto.
- CRINGAN, A.T. 1952. Report on a Trip to Cedar Lake — Perrault Lake-Sioux Lookout District. 6 p. (July 7-11, 1952). Library, Whitney Block, Toronto.
- CRINGAN, A.T. 1956. Preferences for Salt Compounds (Contents of a letter received by A.T. Cringan from York Edwards on this subject). 1 p. Library, Whitney Block, Toronto.
- CRINGAN, A.T. 1956. Some Aspects of the Biology of Caribou and a Study of the Woodland Caribou Range of the Slate Islands, Lake Superior, Ontario. 300 p. (*In* M.A. Thesis, University of Toronto).
- CRINGAN, A.T. 1957. History, Food Habits and Range Requirements of the Woodland Caribou of Continental North America. 17 p. (Repr. from Trans. of 22nd N. Amer. Wildl. Conf., Mar. 4-6, 1957, Washington, D.C., p. 485-501). Library, Whitney Block, Toronto.
- CRINGAN, A.T. 1957. Possible Effects of Forest Fire on Big Game in the Sioux Lookout Forest Protection District. 18 p. (Fish and Wildl. Mgt. Rept. #36, Aug. 1, 1957). Library, Whitney Block, Toronto.
- CUMMING, H.G. 1954. Report on Helicopter Survey of Caribou on Slate Islands. 6 p. (Fish and Wildl. Mgt. Rept. #25, Oct. 1955). Library, Whitney Block, Toronto.
- CUMMING, H.G. 1955. Report on Helicopter survey of Caribou on Slate Islands. (*In* Fish and Wildlife Management Report 25: 23-26).
- CUMMING, H.G. 1956. Report on 1956 Trip to the Slate Islands. 11 p. (Fish and Wildl. Mgt. Rept. #32, Dec. 1, 1956). Library, Whitney Block, Toronto.
- CUMMING, H.G. 1975. Clumping Behaviour and Predation with Special Reference to Caribou. 24 p. (*In* Proceedings of the First International Reindeer and Caribou Symposium, Aug. 9-11, 1972, University of Alaska, Fairbanks, Alaska). Library, Whitney Block, Toronto.
- CUMMING, H.G. 1978. First Progress Report on Research Project Concerning the Effects of Logging on Caribou. p. 24. (*In* Unpubl. Report). School of Forestry, Lakehead University, Thunder Bay, Ontario.
- CUMMING, H.G. 1979. Second Annual Report on Caribou/Logging Project. p. 6. (*In* Unpubl. Report). School of Forestry, Lakehead University, Thunder Bay, Ontario.
- CUMMING, H.G. 1981. Caribou Country. p. 27-30. *In* Seasons. Summer 1981.
- CUMMING, H.G. 1982. Progress Report in Caribou Research. p. 8. (*In* Unpubl. Report 1980-82). School of Forestry, Lakehead University, Thunder Bay, Ontario.
- CUMMING, H.G. and D.B. BEANGE. 1985. Dispersion and movements of Woodland Caribou near Lake Nipigon in Ontario. J. Wild. Manage. 51(1): 69-79.
- CUSSON, M.R. 1964. Kapuskasing District Annual Report — Status of Fur-bearing, Game Animals and Birds. (*In* Wildlife Management District Annual Reports 1963-64. Part I). Library, Whitney Block, Toronto.

- DALTON, W.J. 1985. Background Report for the Slate Island Provincial Park Management Plan. Unpubl. Report, Terrace Bay District.
- DARBY, W.R. and L.S. DUQUETTE. 1986. Woodland Caribou and forestry in Northern Ontario, Canada. In: Gunn, A., Miller, F.L. and Skjenneberg S. (eds.). Proc. Fourth Inter. Reindeer/Caribou Symp. Rangifer Spec. Issue No.1: 87-93.
- DARBY, W.R. and W.O. PRUITT, Jr. 1984. Habitat use movements and grouping behaviour of Woodland Caribou, *Rangifer tarandus caribou*, in southeastern Manitoba. Can. Field Naturalist. 98(2): 184-190.
- DAUPHINE, T.C., Jr. 1981. Progress Report for Radio Tracking of Caribou. p. 137-161. (In J. Tierney (ed.) Caribou Surveys in Pukaskwa National Park 1972-1979). Parks Canada, Pukaskwa National Park, Marathon, Ontario
- DAWSON, F.N. and D.A. PAYNE. 1985. The 1984-85 aerial survey of Moose and Woodland Caribou in W.M.U. 26. Ont. Min. Nat. Res. unpubl. rep., Cochrane District. 33 p.
- DE VOS, A. 1948. Preliminary Report on the Winter Food Habits of Woodland Caribou, 1948. 2 p. Library, Whitney Block, Toronto.
- DE VOS, A. 1948. Status of the Woodland Caribou in Ontario. 6 p. (Sylva, Vol. 4(1): 17-23). Library, Whitney Block, Toronto.
- DE VOS, A. 1948. The Former and Present Distribution of the Woodland Caribou (*Rangifer caribou*) in Northern Ontario, with Suggestions for its Management. 3 p. (Pres. at 10th Midwest Wildl. Conf., Ann Arbor, Mich., Dec. 9-11, 1948). Library, Whitney Block, Toronto.
- DE VOS, A. 1948. The Woodland Caribou Approaches Extinction. 5 p. Library, Whitney Block, Toronto.
- DE VOS, A. 1949. Report on the Woodland Caribou Crown Game Preserve in Sioux Lookout District. 7 p. Library, Whitney Block, Toronto.
- DE VOS, A. 1949. The Outlook for Woodland Caribou Research in Ontario. 5 p. Library, Whitney Block, Toronto.
- DE VOS, A. 1950. Report on a Preliminary Survey of the Goulais River — Ranger Lake Crown Game Preserve, July 22-July 31, 1950. 13 p. Library, Whitney Block, Toronto.
- DE VOS, A. 1959. Progress in Wildlife Management in Canada. 4 p. (In Int'l Union of Game Biologists; Transactions of the 4th Congress, Sept. 28-Oct. 5, 1959). Library, Whitney Block, Toronto.
- DE VOS, A. Undated. Differences between Moose and Caribou (drawings). 1 p. Library, Whitney Block, Toronto.
- DE VOS, A. Undated. Status of the Woodland Caribou in Ontario. 4 p. (Repr. from Sylva, Vol. 4(1): 17-23). Library, Whitney Block, Toronto.
- DE VOS, A. and R.L. PETERSON. 1951. A Review of the Status of the Woodland Caribou (*Rangifer caribou*) in Ontario. 9 p. (Repr. from J. Mamm., Vol 32(3): 329-337). Library, Whitney Block, Toronto.
- DOUGLAS, C.W. 1954. Proposed Woodland Caribou Capture and Transfer from Slate Islands. 3 p. Library, Whitney Block, Toronto.
- DOUGLAS, C.W. 1955. Slate Island Caribou Investigation, Sept. 12-17, 1955. 10 p. (Fish and Wildl. Mgt. Rept. #31, Oct. 31, 1956). Library, Whitney Block, Toronto.
- DYMOND, J.R. 1949. Wildlife Research in Ontario. 6 p. Library, Whitney Block, Toronto.
- EASON, G. 1982. Caribou Management Proposal, Wawa District. Unpubl. Report, Wawa District. 11 p.
- EASON, G. 1984. Caribou Transfer Proposal, Montreal Island. Unpubl. Report, Wawa District. 18 p.
- ENGLISH, G. 1974. Aerial caribou survey — 1974 Irregular Lake Study Area. Unpubl. Report, Red Lake District. 3 p.
- ENGLISH, G. 1976. Aerial caribou survey — 1976 Irregular Lake Study Area. Unpubl. Report, Red Lake District. 2 p.
- EULER, D.L., B. SNIDER, and H.R. TIMMERMAN. 1976. Woodland Caribou and Plant Communities on the Slate Islands, Lake Superior. 5 p. (In Can Field. Nat. 90(1): 17-21). Library, Whitney Block, Toronto.
- FERGUSON, S.H. 1979. Population dynamics of Pic Island caribou. Unpubl. Report, Terrace Bay District. 18 p.
- FERGUSON, S.H. 1982. Why are caribou on Pic Island? 171 p. (In M.A. Thesis. Univ. of Victoria).
- FERGUSON, S.H. and R.S. FERGUSON. 1980. Typed report for Parks Canada. (Pukaskwa National Park, Marathon, Ontario). 16 p.
- FERGUSON, S.H. and R.S. FERGUSON. 1980. On the presence and uniqueness of Pic Island caribou. Unpubl. Report, Terrace Bay District. 23 p.
- FERGUSON, S.H. and R.S. FERGUSON. 1981. The island caribou. Unpubl. Report, Terrace Bay District. 50 p.



- FERGUSON, S.H. and R.S. FERGUSON. 1983. Pic Island caribou: Annual Report 1982. Unpubl. Report, Terrace Bay District. 31 p.
- FISCHER, C.A., G.L. FISHER, G.H. KLASSEN and D.C. THOMPSON. 1977. 75 p. Studies of moose and woodland caribou in Northeastern Manitoba, and Northwestern Ontario. Renewable Resources Consulting Services Ltd. for the Polar Gas project.
- FRUETEL, M.J. 1987. The biology of the gastro-intestinal helminthes of woodland and barren-ground caribou (*Rangifer tarandus*). MSc. Thesis, Lakehead Univ., Thunder Bay, 114 p.
- GAGNON, A. 1959. Trappers Meeting, Fort Albany-Cochrane Interim Reports. 4 p. (*In Wildlife Management District Annual Reports 1958-59. Part I*). Library, Whitney Block, Toronto.
- GAGNON, A. 1960. Caribou Survey, Patricia East, 1960. 3 p. Library, Whitney Block, Toronto.
- GAGNON, A. 1960. Caribou Survey Report — Cochrane District Interim Reports. 6 p. (*In Wildlife Management District Annual Reports 1959-60. Part I*). Library, Whitney Block, Toronto.
- GIBBARD, H.J. 1959. Patricia East, Caribou Survey, 1959. 3 p. Library, Whitney Block, Toronto.
- GIBSON, B.H. 1962. Winter Aerial Census of Woodland Caribou (*Rangifer tarandus*) in the Pickle Crow-Armstrong and Kowkash-Martin Falls Regions. 32 p. Library, Whitney Block, Toronto.
- GISON, B.H. 1970. Geraldton Forest District fish and wildlife management plan for the period 1970-1980 inclusive. (*In Unpub. Report, OMNR Geraldton District*). 95 p.
- GLADSTONE, B. 1988. Carry On Caribou. Landmarks 6(1):6-9.
- GODDARD, J. 1959. Caribou (*Rangifer tarandus*) Survey of the Slate Islands of Lake Superior, May-June 1959. 5 p. Library, Whitney Block, Toronto.
- GODDARD, J. 1961. Aerial Census of Woodland Caribou (*Rangifer tarandus*) in the Attawapiskat-Wunnummin Region, 1961. 7 p. (Fish and Wildl. Review Vol 1(3): 8-14). Library, Whitney Block, Toronto.
- GODDARD, J. 1961. Report of an Aerial Census of Woodland Caribou (*Rangifer tarandus*) Populations in the Attawapiskat-Wunnummin Region During the Winter of 1961. 105 p. Library, Whitney Block, Toronto.
- GOLLAT, R. 1976. The fish and wildlife values of the Black Bay peninsula and offshore Lake Superior Islands, a preliminary investigation. Unpubl. Report, Nipigon District. 103 p.
- GOLLAT, R. 1976. Preliminary investigations of the fish and wildlife values on the islands of Lake Nipigon. 99 p. Unpubl. Report, Nipigon District. 99pp.
- GORSLINE, T. 1978. The caribou crisis. *In Ontario Out of Doors*, April. 78 p.
- GOW, J. 1966. Big Game Killed by Means other than Legal Hunting — Geraldton District, 1966. 2 p. Library, Whitney Block, Toronto.
- GOW, J. 1967. Geraldton District Fish and Wildlife Annual Report — 1967. 1 p. (*In Wildlife Management District Annual Reports 1967-68. Part I*). Library, Whitney Block, Toronto.
- GOW, J. 1968. Big Game Killed by Means other than Legal Hunting — 1968, Geraldton District. 2 p. Library, Whitney Block, Toronto.
- GRAY, P.A. 1978. Native Peoples Harvest of Woodland Caribou (*Rangifer tarandus caribou*) in the West Patricia Planning Area, 1960 to 1978. 39 p. Library, Whitney Block, Toronto.
- GREET, R.M. 1980. A study of mid-winter behaviour patterns of captive woodland caribou with special reference to shelter seeking behavior. 74 p. (*In Unpubl. Report. School of Forestry, Lakehead Univ., Thunder Bay, Ontario*).
- HAGAN, D. 1968. Aerial Caribou Survey — Irregular Lake Study Area, 1968. 6 p. Library, Whitney Block, Toronto.
- HAGAN, D. and L. PENNEY. 1969. Fish and Wildlife Management Plan, Sioux Lookout District. 89 p. Library, Whitney Block, Toronto.
- HALL, F.L. 1958. Polar Bear and Caribou Observations Near Moosonee, Nov. 13, 1958. 2 p. Library, Whitney Block, Toronto.
- HAMILTON, G.D. 1978. Aerial Census of Woodland Caribou in the West Patricia Planning Area. 14 p. (West Patricia Land Use Plan, Wildl. Tech. Rept. No. 6, December 1978). Library, Whitney Block, Toronto.
- HAMILTON, G.D. 1978. Woodland Caribou: A Review with Special Reference to the West Patricia Planning Area. 39 p. (West Patricia Land Use Plan, Wildl. Tech. Rept. No. 4). Library, Whitney Block, Toronto.

- HAMILTON, G.D. 1978. Woodland caribou: A Review with Special reference to West Patricia Planning Area. West Patricia Land Use Plan Tech. Report No.4. Unpubl. Report, Kenora District.
- HAMILTON, G.D. 1979. A Study of Some Woodland Caribou Wintering Areas Within The West Patricia Planning Area. 21 p. (West Patricia Land Use Plan, Wildl. Tech. Report No. 16). Library, Whitney Block, Toronto.
- HAMILTON, G.D. 1979. Reassessment of Woodland Caribou Populations in the West Patricia Planning Area. 9 p. (*In* Wildlife Technical Report No. 15. Aug. 1979). Library, Whitney Block, Toronto.
- HAMILTON, G.D. 1980. Moose and Woodland Caribou in West Patricia. 4 p. (Ont. Fish and Wildl. Review Vol. 19(1): 7-10). Library, Whitney Block, Toronto.
- HAMILTON, G.D. 1984. Native Use of Moose and Woodland Caribou in the Cat Lake Band Area, northwestern Ontario. Masters Thesis, Lakehead University Library. 113 p.
- HANSSON, C. 1963. Big Game Animals Killed by Means other than Legal Hunting, Kenora District — 1963. 6 p. Library, Whitney Block, Toronto.
- HANSSON, C. 1967. Big Game Animals Killed by Means other than Legal Hunting in the Kenora District — 1967. 8 p. Library, Whitney Block, Toronto.
- HANSSON, C. 1967. The Present Status of Woodland Caribou in the Kenora District — 1966-67. 6 p. Library, Whitney Block, Toronto.
- HARKNESS, W.J.K. 1958. Fish and Wildlife Ontario — 1958, An Establishment and Working Plan. 36 p. Library, Whitney Block, Toronto.
- HEPBURN, R. and D.W. SIMKIN. 1960. Patricia Resources Inventory — Caribou Project, 1960. 3 p. Library, Whitney Block, Toronto.
- JACKSON, G. 1981. Woodland caribou: report prepared for Geraldton District land use plan. Unpubl. Report, Geraldton District, 13 p.
- KERR, A.J. 1947. Report on the Comparative Moose and Caribou Kill in the Southern James Bay the Winters 1945-46 and 1946-47. Library, Whitney Block, Toronto.
- LANKESTER, M.W. 1976. A protostrongylid nematode of woodland caribou and its implications in moose management. Proc. N. Am. Moose Conf. Workshop 12: 173-190.
- LANKESTER, M.W., V.J. CRICHTON and H.R. TIMMERMAN. 1976. A protostrongylid nematode (Strongylida: Protostrongylidae) in woodland caribou (*Rangifer caribou*). Can. J. Zool. 54: 680-684.
- LANKESTER, M.W. and M. FRUETEL. 1986. 16 p. Pathogenic helminths of Woodland Caribou. (*In* Renewable Resources Research Seminar, Jan. 30-31, 1986, Toronto).
- LOCKWOOD, J.W. 1957. Annual Report — White River District — Game and Fur Animal Abundance. 8 p. (*In* Wildlife Management District Annual Reports 1956-57. Part II). Library, Whitney Block, Toronto.
- LUCKING, E. 1979. Woodland caribou: a compendium of sightings in Kapuskasing District. Unpubl. Report, Kapuskasing District. 5 p.
- LUMSDEN, H.G. 1955. Caribou Inventory, 1955-56. 1 p. Library, Whitney Block, Toronto.
- MCGUIRE, S. 1988. Caribou cowboys have a gentle touch. Landmarks, 6(1): 10-11.
- MECH, L.D., M.E. NELSON and H.F. DRABIK. 1982. (*In* Reoccurrence of caribou in Minnesota. Am. Midl. Nat. 108(1): 206-208).
- MULLEN, C.E. 1981. Preliminary report, Woodland Caribou survey, Aril-May 1973. 15-20 p. (*In* J. Tierney (ed.) Caribou surveys in Pukaskwa National Park, 1972-1979. Parks Canada, Pukaskwa National Park, Marathon, Ontario).
- PAULSEN, H.E. 1975. Caribou Studies. 20 p. (*In* Fish and Wildlife Nipigon District Annual Report 1974/75. Ed. P.B. Hogan. p. 122-134). Library, Whitney Block, Toronto.
- PAULSEN, H.E. 1976. 1975-1976 Caribou Studies. 2 p. (*In* Nipigon District Annual Fish and Wildlife Report 1975/76. Ed. R.J. Swanson. p. 128-129). Library, Whitney Block, Toronto.
- PAULSEN, H.E. 1976. Winter Range of the Armstrong Caribou Herd. 3 p. (*In* Nipigon District Annual Fish and Wildlife Rept. 1975/76. Ed. R.J. Swanson. p. 124-126). Library, Whitney Block, Toronto.
- PAULSEN, H. 1984. Caribou lichen plots. 76-77 p. (*In* P. Burkhardt (ed.) Nipigon District Annual Fish and Wildlife Report).
- PERRIE, C.E. 1954. Report on Aerial Survey of Slate Islands — January 26th, 1954. 2 p. (Fish and Wildl. Mgt. Rept. #31, Oct.1, 1956). Library, Whitney Block, Toronto.




- PETERSON, R.L. 1966. The mammals of Eastern Canada. 465 p. Oxford University Press, Toronto.
- SCHRAEDER, H. 1982. A Discussion on the effects of logging on woodland caribou (*Rangifer tarandus caribou*) with special inference to the Armstrong airport herd. Unpubl. Report, Nipigon District. 22 p.
- SHANNON, J.A. 1957. Big Game Status of the Woodland Caribou Crown Game Preserve. 2 p. Library, Whitney Block, Toronto.
- SHIELDS, W.F. 1959. Patricia East Caribou Survey. 11 p. (Sylva, May -June 1959, p. 4-13). Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1958. Woodland Caribou Investigations in West Patricia, 1958. 6 p. (Pres. at the Fort Frances Fur Advisory Comm. Meeting, Apr. 10-11, 1958). Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1959. Future Utilization and Management of Caribou in the Patricias. 10 p. (Pres. at the Port Arthur Fur Advisory Comm. Meeting, Apr. 23-24, 1959). Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1959. Survey of Caribou in Woodland Caribou Crown Game Preserve, 1959. 2 p. Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1960. Caribou Hunter Interviews in Patricia Central and Patricia West, 1959. 8 p. (Fish and Wildl. Mgt. Rept. #51, April 1960). Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1960. Woodland Caribou Project, July 1959. 10 p. Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1960. Woodland Caribou Project, Spring 1960. 11 p. Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1960. Woodland Caribou Project, Summer 1960. 20 p. (Resource Mgt. Rept. #60, November 1961). Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1961. Aerial survey for caribou in Hudson Bay Lowland. Unpubl. Report, Geraldton District.
- SIMKIN, D.W. 1961. Spring Caribou Calving Study, 1961. 3 p. Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1965. Aerial survey for caribou — Irregular Lake study area. Ontario Department of Lands and Forests. Unpubl. Res. Report.
- SIMKIN, D.W. 1965. Preliminary Report of the Woodland Caribou Study in Ontario. Ontario Department of Lands and Forests Sect. Report (Wildl.) No. 59. 75 p. Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1965. Woodland Caribou in Ontario. 7 p. (Fish and Wildl. Review Vol. 4(4): 2-8). (2 copies) Library, Whitney Block, Toronto.
- SIMKIN, D.W. 1966. Aerial survey for caribou in area northwest of Big Trout Lake, Feb. 1966. Ontario Department of Lands and Forests, Unpubl. Res. Report, Maple. 9 p.
- SIMKIN, D.W. 1967. Aerial survey for caribou between Severn and Winisk rivers, March 1967. Ontario Department of Lands and Forests, Unpubl. Res. Report, Maple. 5 p.
- SNIDER, J.B. 1977. Caribou rutting behavior on the Slate Islands. Unpubl. Report, Terrace Bay District. 2 p.
- SNIDER, J.B. 1977. Slate Island expedition of May 30th, 1977 to June 5th, 1977. Unpubl. Report, Terrace Bay District. 7 p.
- SNIDER, J.B. 1983. Notes on the Caribou of the Slate Islands: March Trip 1983. Library, Whitney Block, Toronto. 12 p.
- SNIDER, J.B. 1986. Slate Island investigations of March 1985: the year of the die-off. Unpubl. Report, Terrace Bay District. 8 p.
- SNIDER, J.B. 1987. Slate Islands investigations of April, 1986. Unpubl. Report, Terrace Bay District. 3 p.
- SNIDER, J.B. and S.H. FERGUSON. 1978. Wolf scat analysis, Coldwell Peninsula. Unpubl. Report, Terrace Bay District. 1 p.
- SNYDER, L.L. 1938. A faunal investigation of western Rainy River district Ontario. Trans. Roy. Can. Inst. Vol. 14, Part 1: 157-181, Toronto
- SNYDER, L.L., E.B.S. LOGIER and T.B. KURATA. 1942. A faunal investigation of the Sault Ste. Marie region, Ontario. Contr. Roy. Ont. Mus. Zool. No. 21.
- STEWART, R.W. 1977. The 1976-77 Caribou Aerial Inventory of WMU 26. Ont. Min. Nat. Res. unpubl. rep., Cochrane District.
- ST. JULES, S. 1965. Status of Fur-bearing, Game Animals and Birds — Kapuskasing District. (In Wildlife Management District Annual Reports 1964-65. Part I). Library, Whitney Block, Toronto.
- ST. JULES, S. 1966. Indian Kill of Big Game — Kapuskasing District. 1 p. (In Wildlife Management District Annual Reports 1965-66. Part I). Library, Whitney Block, Toronto.



- THOMPSON, J.E. 1986. Population and harvest surveys for the Ontario Hudson's Bay Lowland caribou. Unpubl. Report, Moosonee District 36 p.
- TIMMERMAN, H.R., J.B. SNIDER and D. EULER. 1974. The Slate Islands. Unpubl. Report, Terrace Bay District. 41 p.
- TIMMERMAN, H.R. 1985. Caribou relocation project, St. Ignace Island, Nipigon District, North Central Region. Unpubl. Report, North Central Region, 13 p.
- VARRIN, P.W. 1976. Caribou and Moose Surveys and Tagging — Lake Nipigon Islands — Summer 1976. 9 p. (*In* Nipigon District Annual Fish and Wildl. Rept. 1976/77. Ed. P.W. Varrin. p. 41-49). Library, Whitney Block, Toronto.
- VARRIN, P.W. 1976. Caribou Studies — South Half of Lake Nipigon — 1976. 2 p. (*In* Nipigon District Annual Fish and Wildlife Rept. 1976/77. Ed. P.W. Varrin. p. 50-51). Library, Whitney Block, Toronto.
- VARRIN, P.W. 1977. Caribou Aerial Surveys — 1977. 3 p. (*In* Nipigon District Annual Fish and Wildlife Rept. 1976/77. Ed. P.W. Varrin. p. 71-73). Library, Whitney Block, Toronto.
- VARRIN, P.W. 1978. The 1977 Caribou-Moose Tagging Report and Winter '78 Aerial Caribou Census. 12 p. (*In* Nipigon District Annual Fish and Wildlife Report 1977-78. Ed. P. Burkhardt. p. 129-139). Library, Whitney Block, Toronto.
- VARRIN, P.W. 1979. Reassessment of Woodland Caribou Populations in the West Patricia Planning Area. 9 p. (Wildl. Tech. Rept. No. 15. Aug. 1979). Library, Whitney Block, Toronto.
- WALROTH, R. 1980. Caribou field trip, Whitesand River area, September 25, 1979; Winter observations of Armstrong caribou herd 1979-80; Assessment of caribou lichen plots. (*In* Ont. Min. Nat. Res. Nipigon District Annual Fish and Wildlife Report 1979/80. p. 169-175).
- WALROTH, B. 1981. Caribou pellet group and lichen (*Cladonia*) survey, Armstrong, May 1981. (*In* P. Burkhardt (ed.) OMNR Nipigon District Annual Fish and Wildlife Report 1982-83. p. 135-148).
- WALROTH, R. 1982. Caribou pellet group and lichen *Cladonia* survey, Armstrong, May 1982. Unpubl. Report, Nipigon District. 16 p.
- WEPRUK, R. 1986. Woodland caribou investigations of Woodland Caribou Park, 1985, draft. Woodland Caribou Provincial Park Technical Report No. 3., Unpubl. Report, Red Lake District, 41 p.
- WILD, J. 1981. The use of conifer shelter by captive woodland caribou (*Rangifer tarandus caribou* Gmelin) in winter. 65 p. (*In* Unpubl. Report). School of Forestry, Lakehead University, Thunder Bay.
- WYETT, W. and G. KEESEY. 1978. Faunal investigations in Pukaskwa National Park. 38 p. (*In* Unpubl. Report Pukaskwa National Park, Marathon, Ontario).





4175  
(1 k P.R., 89 01 30)  
ISBN 0-7729-4449-0